



Anatomy of a Duality

Clifford V. Johnson[†]

*Department of Physics and Astronomy
University of Kentucky
Lexington, KY 40506-0055 USA*

Abstract

The nature of M-theory on $K3 \times \mathcal{I}$, where $\mathcal{I}$ is a line interval, is considered, with a view towards formulating a “matrix theory” representation of that situation. Various limits of this compactification of M-theory yield a number of well known $\mathcal{N}=1$ six dimensional compactifications of the heterotic and type I string theories. Geometrical relations between these limits give rise to string/string dualities between some of these compactifications. At a special point in the moduli space of compactifications, this motivates a partial definition of the matrix theory representation of the M-theory on $K3 \times \mathcal{I}$ as the large N limit of a certain type IA orientifold model probed by a conglomerate of N D-branes. Such a definition in terms of D-branes and orientifold planes is suggestive, but necessarily incomplete, due to the low amount of supersymmetry. It is proposed —following hints from the orientifold model— that the complete matrix theory representation of the $K3 \times \mathcal{I}$ compactified M-theory is given by the large N limit of compactification —on a suitable “dual” surface— of the “little heterotic string” $\mathcal{N}=1$ six dimensional quantum theories.

11th November 1997

email: [†]cvj@pa.uky.edu

Contents

1. Motivation and Background	3
1.1. <i>Compactifying M–Theory</i>	3
1.2. <i>The DMW Point: Heterotic/Heterotic Duality</i>	4
1.3. <i>Relation to the GP Model</i>	4
1.4. <i>Searching for a Matrix Theory Representation</i>	6
1.5. <i>Outline</i>	7
2. Some Orientifold Models	8
2.1. <i>The GP Orientifold Model: Type IB on K3</i>	8
2.2. <i>Another Orientifold Model: Type IA on K3</i>	10
2.3. <i>M–Theory on $K3 \times \mathcal{I}$</i>	12
2.3.1. <i>Digression on D8–Branes and O8–Planes</i>	12
2.3.2. <i>Inclusion of K3, D4–Branes and O4–Planes</i>	13
2.4. <i>The $E_8 \times E_8$ and $Spin(32)/\mathbb{Z}_2$ Heterotic Strings on K3</i>	15
2.5. <i>Geometrical Picture of the Dualities</i>	16
2.6. <i>Origin of Heterotic/Heterotic Duality</i>	17
2.7. <i>Heterotic/Heterotic Duality as Eleven Dimensional Electromagnetic Duality</i>	21
3. Probing with D–Branes	24
3.1. <i>Probing with D1–Branes: The Model</i>	25
3.1.1. <i>Supersymmetry</i>	25
3.1.2. <i>The 1–1 Strings</i>	26
3.1.3. <i>The 1–9 Strings</i>	28
3.1.4. <i>The 1–5 Strings</i>	29
3.1.5. <i>The 9–9, 5–5 and 5–9 Strings</i>	29
3.1.6. <i>Two Puzzles, and Their Solution</i>	30
3.1.7. <i>The Closed Strings</i>	33
3.2. <i>Probing with D1–Branes: The Geometry</i>	35
3.2.1. <i>Instantons, ALE Spaces, HyperKähler Quotients and ADHM Data</i>	35
3.2.2. <i>The Moduli Space of GP Models and Probe Models</i>	37

3.2.3. <i>The Special Point</i>	40
3.3. <i>Probing with D0-branes and D5-branes</i>	42
3.3.1. <i>D0-Brane Probes</i>	42
3.3.2. <i>D5-Brane Probes</i>	42
4. Some (Partial) Matrix Theory Proposals	43
4.1. <i>Matrix Theory of M-theory on $X \times \mathcal{I}$</i>	44
4.2. <i>Matrix Theory of the $E_8 \times E_8$ Heterotic String on $X \times \mathcal{I}$</i>	44
4.1. <i>Matrix Theory of M-theory on $K3 \times \mathcal{I}$ and its Heterotic Limits</i>	45
5. Beyond the Special Point	47
6. Discussion and Conclusion	47
6.1. <i>Summary</i>	47
6.2. <i>New Directions</i>	47
Acknowledgements	50
References	51

1. Motivation and Background

1.1. Compactifying M-Theory

Consider the case of M-theory compactified on the surface $K3 \times \mathcal{I}$, where $\mathcal{I}$ is a line interval of length L . Consistent compactification[1] requires the introduction of 24 M5-branes, transverse to the $K3 \times \mathcal{I}$, in order to cancel the resulting chiral anomaly in the six dimensional transverse space $\mathbb{R}^6$. The anomaly owes its presence to essentially the curvature of $K3$, which has Pontryagin number 24 in the appropriate units. Put another way, $K3$'s curvature produces 24 units of magnetic $A^{(3)}$ charge in the background, which may be canceled by the 24 units of magnetic charge of the M5-branes[2], M-theory's basic dynamical magnetic $A^{(3)}$ charge carriers at low energy¹.

As is now standard lore, in the Hořava–Witten limit ($L \rightarrow 0$), this compactification of M-theory yields a discrete *family* of compactifications of the $E_8 \times E_8$ heterotic string on $K3$. The resulting (ten dimensional) heterotic string coupling is related to the interval size as $\lambda \sim L^{3/2}$.

There are two distinct categories of heterotic vacua obtained in this way. The first is “ordinary” heterotic string vacua, where all of the massless spectrum can be accounted for in string perturbation theory. Choices made for the distribution of M5-branes translate directly into the choice of embedding n_1 of the instantons in one E_8 and the other $n_2 = 24 - n_1$ into the second E_8 . This corresponds, in M-theory, to the partitioning of the transverse M5-branes such that there are n_1 at one endpoint of the interval $\mathcal{I}$, and n_2 at the other, before taking the heterotic limit.

There are also heterotic string vacua with n_T massless states in their spectrum which transform as tensors. The chiral anomaly can allow them as long as $n_1 + n_2 + n_T = 24$. These are not to be understood as completely perturbative heterotic vacua, but ones which contain contributions from non-perturbative effects which make their presence felt all the way down to the weak coupling limit. In M-theory ($L \neq 0$), this seemingly mysterious statement is realized explicitly in terms of making the choice (n_1, n_2) for the numbers of M5-branes at the respective ends of the interval, while soaking up the rest of $K3$'s curvature-induced magnetic $A^{(3)}$ charge with n_T M5-branes in the interior. The tensor massless degrees of freedom living one on each M5-brane world volume, are now free to contribute to the overall spectrum. In taking the stringy limit, that arrangement yields this distinct second class of heterotic string vacua.

In both cases, the heterotic string itself arises from wrapping the M2-brane[3] on the interval $\mathcal{I}$, the boundary of the brane giving a string in the transverse space. M-theory details completely independent of L (*i.e.*, residing on the ends of the interval where the

¹ Here, $A^{(3)}$ denotes the three-form potential of eleven-dimensional supergravity, the low energy limit of M-theory.

boundary of the M2-brane lives) like the instantons, become identifiable as perturbative string physics in the $L \rightarrow 0$ limit. Meanwhile, M-theory physics of the bulk (which is also the bulk of the M2-brane) is not perturbatively described in the limit.

1.2. The DMW Point: Heterotic/Heterotic Duality

Turning back to the $n_T=0$ cases, it is known[4] that there is one special case where there exists a different heterotic limit. One first performs an eleven dimensional electromagnetic duality operation with respect to $A^{(3)}$. This necessarily involves the exchange of the M2-brane with an M5-brane oriented with four of its directions on $K3$, and one in the $\mathbb{R}^6$. (It therefore contributes nothing to the anomaly mentioned earlier.) Sending the volume of the $K3$ to zero results in a string in $\mathbb{R}^6$ which is the heterotic string.

There is actually only one way to do this geometrically[4], and this is for choice (12,12) distribution of the transverse M5-branes. Only then can the $K3$ be seen to support no residual magnetic charge in bulk and therefore be consistently shrunk away to zero with an M5-brane wrapped on it².

That is the eleven dimensional geometrical origin of heterotic/heterotic duality in six dimensions, defining the special Duff–Minasian–Witten point[4] in the moduli space of $E_8 \times E_8$ heterotic string $K3$ compactifications. This is the (long anticipated[5]) $\mathcal{N}=1$ string/string duality in six dimensions which is a natural ancestor of $\mathcal{N}=2$ string duality in four dimensions.

1.3. Relation to the GP Model

Fortuitously, there exists yet another duality (discussed by Berkooz, Leigh, Polchinski, Schwarz, Seiberg and Witten[6]) which relates the special (DMW) point to a type IIB orientifold[7][8] model representing the type IB string compactified on $K3$, called the Gimon–Polchinski model[9]³.

There are a number of important points to be made here about the establishment of this duality. More correctly, the heterotic/heterotic DMW point is dual to precisely that special point in the moduli space of GP models where there might be some hope to make strong/weak coupling duality contact with a perturbative heterotic string compactified

² Really, only a partial argument was presented in ref.[4]; It was not clear there how the $K3$ in the resulting heterotic compactification reappears. This M-theory argument will be completed in this paper.

³ A special point in the moduli space of the Gimon–Polchinski (GP) models was also recognized as a consistent spectrum by Bianchi and Sagnotti in ref.[10]. This is the point with largest possible gauge group $U(16) \times U(16)$. In the interests of historical accuracy, we might call this point the BS–GP point, reserving the name GP for the more general class of models found in ref.[9].

on $K3$. (This point, which we'll call the "special GP model" has no regions of spacetime where the dilaton has a gradient, thereby promising a heterotic conformal field theory dual description. Such orbifold conformal field theories were presented in refs.[11,6].) One might have anticipated such a duality as a descendent of the ten dimensional strong/weak coupling duality between the $SO(32)$ type IB string and the $SO(32)$ heterotic string. However, there is no straightforward geometrical reason why such a heterotic dual to the GP model should have anything to do with heterotic/heterotic duality: The ten dimensionally motivated expectation should only generically predict duality to some $K3$ compactification of the $SO(32)$ heterotic string.

Part of the work of ref.[6] in showing that the special GP model was indeed dual to the DMW model involved proving that a much more complicated example of a phenomenon familiar in toroidal heterotic compactifications occurs here: There is a T-duality between the $(12,12)$ $K3$ compactified $E_8 \times E_8$ heterotic string and a $K3$ compactification of the $SO(32)$ heterotic string. In order to make this correspondence, it has to be realized that in the latter case, special choices are made for the instantons which refer to their structure as $Spin(32)/\mathbb{Z}_2$ instantons and not merely $SO(32)$ ones.

Once this T-duality connection is made (a perturbative demonstration which boils down to making careful choices of $Spin(32)/\mathbb{Z}_2$ vector bundles and choosing an appropriate Wilson line), the ten dimensional expectation goes through and makes a connection between the special GP model and the DMW point.

Indeed, there are some amusing facts which bolster the connection[6]: For example, it is easy to construct the string solitons on the type IB side which eventually become dual heterotic strings on the heterotic side. They are simply the two distinct types of D-string probe one can make in the six uncompactified dimensions. One is simply the ordinary D1-brane in $\mathbb{R}^6$, while the other is made by wrapping a D5-brane on the $K3$ to give a D-string in $\mathbb{R}^6$. Note that in the GP model, these two D-string solitons are T-dual to each other under inversion of the volume of the (orbifold) $K3$. Under the strong/weak coupling duality, one of these strings becomes light and becomes the heterotic string, while the other becomes an ordinary heterotic stringy soliton. The Type IB model's $K3$ inversion T-duality symmetry, which exchanges the two D-strings, can now be seen to be "pulled back" under the type IB/heterotic (on $K3$) map to the DMW heterotic/heterotic duality, explaining the presence of a dual heterotic string.

In the first half of this paper, all of these orientifold facts will be explicitly linked to the facts presented in the previous subsection (1.2) by direct connection between the orientifold models and M-theory on one side, and the heterotic models and M-theory on the other side. In the process, a number of puzzling M-theory statements in relation to heterotic/heterotic duality will be clarified and extended.

1.4. Searching for a Matrix Theory Representation

By this point in the discussion, the informed (but patient) reader may wonder why we have gone to all of this trouble to recall the above interesting collection of facts. The goal of the present paper is to try to begin to understand how to construct a matrix theory representation of the $K3 \times \mathcal{I}$ compactified M-theory. It is not clear just how to do this directly from the original Matrix theory prescription[12], which gave rise to a representation of uncompactified M-theory. The technology of compactification on a surface X involves[12,13] studying the large N limit of supersymmetric theories whose target space is a “dual” space $\tilde{X}$. In the case of toroidal compactifications on $X=T^n$ where $n < 4$, the prescription is completely given in terms of $n+1$ dimensional $SU(N)$ Yang–Mills theory, and the “dual” space which will be its target space is simply the torus $\tilde{T}^n$ made by inverting the radii of all of the n constituent circles.

Beyond that class of compactifications, the situation is less clear, principally due to the fact that the ultraviolet behaviour of Yang–Mills theory in higher dimensions is ill-defined. (Also, beyond the case of tori, it is not clear what the “dual” surface should be.) In the case of T^4 , however, it was shown that certain facts about field theory conspire to allow a sensible conjecture[14] about the existence of a sensible Matrix theory definition in terms of the large N limit of a certain six dimensional superconformal fixed point theory. The existence of the appropriate field theory was confirmed in ref.[15], and the Matrix theory consequences, obtained by placing it on $\tilde{T}^5$, were studied further, by testing that known type II string U -duality symmetries are predicted[15].

Even in that case, there is enough supersymmetry and geometrical symmetry of the compactification space (and its dual) to sensibly motivate a complete matrix description within the matrix theory framework. For the cases we would like to consider here, there is less supersymmetry and geometrical symmetry of the target space to help in finding a matrix theory representation.

One way to proceed might be to note that the compactification space which we wish to consider is geometrically realizable (at special points) by orbifolds of tori. Perhaps progress may be made by considering orbifolding the matrix theory definition of M-theory on T^5 [14,15]. The problem with this is of course that we have only a partial understanding of what it means to directly orbifold in either M-theory or matrix theory, and so this is not an avenue of approach without potential wrong turns.

Our starting point is to go back to the spirit of the original matrix theory representation of M-theory and find a well-understood D0-brane background in string theory to endow with an eleven dimensional interpretation, building a matrix theory from the resulting world-volume theory. This approach is not without its pitfalls either. The models we may construct in such a case will of course not have nearly enough supersymmetry to allow the powerful statements about the nature and existence of certain D0-brane bound states at large N —of the type made in the original matrix theory proposal[12]— to be made.

Therefore we only expect a partial definition of the required matrix theory to arise from considering complicated D0-brane backgrounds.

We might hope, however, that we can deduce certain properties of the matrix theory we need by using as many facts as we can, from as many sources as possible. This is the place where the heterotic/heterotic discussion at the beginning comes in. At that special point in the moduli space of M-theory backgrounds, we have a wealth of facts which may help pin down the properties of the representation, (which we might hope is some large N limit of a special type of quantum theory compactified on a “dual” space to the $K3 \times \mathcal{I}$, where “dual” needs to be defined.)

The very existence of the special properties of the DMW point might have been motivation enough to try considering its consequences for a matrix theory proposal⁴. As luck would have it though, we are blessed with a complete D-brane model —more precisely, an orientifold model— which serves as a representation of the special heterotic/heterotic point.

We will therefore pursue our quest to find the required matrix model representation by constructing a variant of the special point of the GP model in the *type IA* theory. The matrix theory representation of the compactification of M-theory on $K3 \times \mathcal{I}$ will be thus partially motivated as the large N limit of a certain type IA orientifold model probed by a conglomerate of N D0-branes, and by extension, also the large N limit of D1- and D5-brane probes of type IB orientifold models.

By the end of the discussion, we will have reason to conjecture that more generally, the complete matrix theory representation of M-theory on $K3 \times \mathcal{I}$ is given by the large N limit of a compactification —on a suitable “dual” surface— of the “little $E_8 \times E_8$ heterotic string” $\mathcal{N}=1$ six dimensional quantum theory.

1.5. Outline

In section 2 we begin by reminding the reader of some of the details of the GP type IIB orientifold model. We then introduce a new (but closely related to the GP model) orientifold model in the type IIA theory, and translate the properties of the GP model into this new model. We then study the strong coupling limit of this type IA model, which is an M-theory $K3 \times \mathcal{I}$ compactification with M-branes. From there we take a weak coupling limit which yields heterotic string theory.

As promised, along the way we will recall, derive, rederive and strengthen our knowledge of properties of the heterotic/heterotic duality, in terms of its image dualities in the various dual contexts of type IB, type IA and M-theory. (The interplay between the latter two is the newly presented context.) By the end of the section, we will have thus firmly recast

⁴ Indeed, this was the question which led the author to embark upon this project.

our knowledge of heterotic/heterotic duality in type IA/M-theory terms, in preparation for section 3.

In section 3, we take our type I models and study them as backgrounds for D-brane propagation. We consider therefore a model of quantum mechanics, for the case of probing local parts of the $K3 \times \mathcal{I}$ geometry with D0-branes, a 1+1 dimensional theory when L_M is small, resulting from D1-brane probes, and later a certain 5 + 1 dimensional quantum theory compactified on a dual surface to $K3 \times \mathcal{I}$ for the case of the full compactification.

In section 4 we submit these models as partial definitions of the Matrix theory representation of M-theory on $K3 \times \mathcal{I}$, at the DMW point, in the infinite momentum frame. As these models inherit the duality properties of the type IA(B) and heterotic models which they were derived from, they are extremely well suited to be used in such a definition. However, due to the lack of enough supersymmetry, there is not a great deal of control over the properties of the short distance information (encoded in the bound state dynamics) which is needed to supplement the model to make it a complete definition.

We discuss such shortcomings of the definition and nevertheless present a proposal for the end result of the inclusion of the missing details in the completion of the models of sections 3 and 4 into a satisfactory definition.

There are good reasons to believe, based on the properties of the orientifold models, that the full 5+1 dimensional theory is the large N limit of the $(0,1)_N$ supersymmetric heterotic theory with $E_8 \times E_8$ global symmetry compactified on a dual surface to $K3 \times \mathcal{I}$. At the DMW point it is also equivalent to a compactification of the $(0,1)_N$ supersymmetric theory with $Spin(32)/\mathbb{Z}_2$ global symmetry. (It is the infinite momentum frame of the M-theory compactification which corresponds to a large N limit of such a configuration.)

Following that identification, it is natural to conjecture (by section 5) that the $E_8 \times E_8$ theory can be used to construct the matrix representation of M-theory on $K3 \times \mathcal{I}$ even away from the DMW point.

2. Some Orientifold Models

2.1. The GP Orientifold Model: Type IB on $K3$

Let us begin by recalling some of the details of the Gimon–Polchinski model[9]. It is a compactification of the type IB string on a $\mathbb{Z}_2$ orbifold of T^4 , realized as the prototype $K3$ orientifold of the type IIB string⁵, where spacetime orbifold symmetries are combined with worldsheet (orientifold) ones. The orientifold model, defined by the group

$$G_\Omega := \{1, R_{6789}, \Omega, \Omega R_{6789}\}, \quad (2.1)$$

⁵ See refs.[16,17] for a wider discussion of $K3$ orientifolds of type IIB.

contains the orbifold symmetry

$$R_{6789} : \{x^6, x^7, x^8, x^9\} \rightarrow \{-x^6, -x^7, -x^8, -x^9\}, \quad (2.2)$$

and the orientifold symmetry $\Omega : z \leftrightarrow \bar{z}$ on the world-sheet which is parameterized by complex coordinate z at string tree level.

The presence of Ω in the group may be thought of as representing a single orientifold plane (“O9-plane”) which fills the whole of spacetime. This plane has -32 units of D9-brane charge. To cancel this charge (as required by “Gauss’ Law” for the Ramond–Ramond potential $A^{(10)}$), 32 D9-branes are introduced.

The presence of ΩR_{6789} represents the introduction of 16 O5-planes, located one at each of the orbifold fixed points of the singular $T^4/\mathbb{Z}_2$ geometry. Their -2 units each of D5-brane charge require 32 D5-branes to be introduced, for consistent $A^{(6)}$ field equations.

In the top part of Table 1 is illustrated the orientations of all of the constituents of the model. The $K3=T^4/\mathbb{Z}_2$ is located in the $\{x^6, x^7, x^8, x^9\}$ directions. The various D5-branes are free to move around in the $K3$, but only in groups of four[9], as there must be a mirror partner of each brane under R_{6789} , and there is an additional pairing for invariance under Ω . We may also introduce Wilson lines to break some of the gauge symmetry of the D5-branes and D9-branes.

type	#	x^0	x^1	x^2	x^3	x^4	x^5	x^6	x^7	x^8	x^9
D9	32	—	—	—	—	—	—	—	—	—	—
O9	1	—	—	—	—	—	—	—	—	—	—
D5	32	—	—	—	—	—	—	•	•	•	•
O5	16	—	—	—	—	—	—	•	•	•	•
D1	N	—	•	•	•	•	—	•	•	•	•
D5'	N	—	•	•	•	•	—	—	—	—	—

Table 1. (Top) Orientation and number of the constituent branes of the Gimon–Polchinski model of the type IB string compactified on a $T^4/\mathbb{Z}_2$ orbifold limit of $K3$ (located in the 6,7,8 and 9 directions). (Bottom) Orientation of branes used to probe the model (see section 3).

The gauge symmetry comes from two sectors, the 9–9 sector and the 5–5 sector, and is maximal when there are no Wilson lines, and the D5-branes are all coincident located

on one of the 16 fixed points. This gives gauge group⁶ $U(16)_5 \times U(16)_9$. (The subscripts denote a gauge group's origin in the 9–9 or 5–5 sector.) 9–9 and 5–5 strings also supply hypermultiplets in the $\mathbf{120} + \overline{\mathbf{120}}$ for each factor, and 9–5 strings supply a $(\mathbf{16}, \mathbf{16})$. The closed string sector supplies 20 neutral hypermultiplets representing the (stringy) geometric deformations of $K3$ ⁷.

As the number of D5–branes which may move around the $K3$ is limited to be a multiple of four, there are a number of distinct models which may be constructed in the GP class which are not connected[9](see a reminder in section 3.2.2). One such model is the special point we are interested in: It has a pair of D5–branes at each O5–plane, thereby canceling the $A^{(6)}$ locally. This unique arrangement is important for the dualities which we want to consider because such a local cancellation in the Ramond–Ramond sector implies a local cancellation in the Neveu–Schwarz–Neveu–Schwarz sector, thus ensuring that the dilaton is constant everywhere. Consequently, any strong/weak dual heterotic model is likely to be realizable as a conformal field theory, as demonstrated explicitly in ref.[6].

For this configuration, there is no gauge group, and there are simply 224 hypermultiplets from the open string sector plus the 20 neutral ones from the closed string sector. (More details on this are to be found in section 3.2.3.)

As pointed out in ref.[9], the four D5–brane dynamical unit constitutes one complete instanton. Therefore there are 8 such instantons in the D5–brane sector, the other 16 coming from the 16 fixed points to make a total of 24, as required. It is crucial to note[6] that these are $Spin(32)/\mathbb{Z}_2$ instantons “without vector structure”, which means that the gauge bundle over the fixed points is chosen in such a way that $U(1)$ instantons can be embedded into $Spin(32)/\mathbb{Z}_2$ and yield the correct Dirac quantization for the adjoint and (say) positive chirality spinor representations, but not for the vector and negative chirality spinor representations. It is $Spin(32)/\mathbb{Z}_2$ heterotic string theory compactified on a $K3$ with such instantons which is T–dual to a $K3$ compactification of the $E_8 \times E_8$ heterotic string, as will be useful for us later. We will also see how these instantons arise naturally in the D–brane probe models in section 3.

2.2. Another Orientifold Model: Type IA on $K3$.

Table 2 shows an arrangement of branes and orientifolds in the type IA theory.

The model has orientifold group

$$G_\Omega = \{1, R_{6789}, \Omega R_5, \Omega R_{56789}\}. \quad (2.3)$$

The compactification is geometrically essentially $K3(\mathbb{Z}_2) \times S^1/\mathbb{Z}_2$ where the S^1 is a compact x^5 of radius $L_{IA}^5/2\pi$. The $\mathbb{Z}_2$ is generated by $R_5 : x^5 \rightarrow -x^5$. The presence of element

⁶ As shown in ref.[6], the $U(16)$'s are broken to $SU(16)$ by one–loop effects. See later.

⁷ This special point is the BSGP point referred to earlier[10][9]

ΩR_5 indicates the presence of two orientifold O8–planes, at the two $\mathbb{Z}_2$ fixed points of the orbifolded circle, forming an interval $\mathcal{I}$. They have charge -16 each in D8–brane units and therefore there are 32 D8–branes for the model to be consistent.

Meanwhile ΩR_{56789} requires 32 O4–planes, split as 16×2 . The sixteen are associated with the fixed points of the $T^4/\mathbb{Z}_2$ $K3$ orbifold, and there are two copies of this arrangement, one for each of the x^5 fixed points, the ends of the interval $\mathcal{I}$.

Each O4–plane has -1 units of D4–brane charge and so there are 32 D4–branes in the model for consistency. They are localized in the five dimensional $K3 \times \mathcal{I}$ space.

type	#	x^0	x^1	x^2	x^3	x^4	x^5	x^6	x^7	x^8	x^9
D8	32	—	—	—	—	—	•	—	—	—	—
O8	2	—	—	—	—	—	•	—	—	—	—
D4	32	—	—	—	—	—	•	•	•	•	•
O4	32	—	—	—	—	—	•	•	•	•	•
D0	N	—	•	•	•	•	•	•	•	•	•

Table 2. (Top) Number and configuration of constituent branes in an orientifold model of the type IA string compactified on a $T^4/\mathbb{Z}_2$ orbifold limit of $K3$ (located in the 6, 7, 8 and 9 directions) times the line interval $S^1/\mathbb{Z}_2$ (located in the 5 direction). (Bottom) Orientation of D0–brane used to probe the model locally realizing a model of matrix quantum mechanics (see section 3).

It should be straightforward to see that this model is essentially T_5 –dual to the GP model compactified on an x^5 circle of radius $L_{\text{IB}}^5/2\pi$, where the relation between parameters of the two models is⁸:

$$\begin{aligned}
L_{\text{IB}}^5 &= \frac{1}{L_{\text{IA}}^5} \\
\lambda_{\text{IB}} &= \frac{\lambda_{\text{IA}}}{L_{\text{IA}}^5} \\
R_{\text{IB}} &= R_{\text{IA}},
\end{aligned} \tag{2.4}$$

where $\lambda_{\text{IA(B)}}$ denotes the string coupling and $R_{\text{IA(B)}}$ denotes the radius of the $K3$ manifold in the type IA(B) theory.

⁸ Here and in what follows, we are not concerned with factors of α' ; nor are we careful about factors of 2π . These will often be omitted.

2.3. M-Theory on $K3 \times \mathcal{I}$

Just as in the case of the GP model, there is a particularly interesting unique configuration where all of the R-R charges are completely canceled locally in the model. The required arrangement is the placement of 16 of the D8-branes on an O8-plane at one end of the interval and the remaining 16 on the O8-plane at the other end[18]. Meanwhile the D4-branes are distributed one per O4-plane, resulting in 16 at each of the copies of $K3$ at the end of the interval.

2.3.1 Digression on D8-branes and O8-planes

In the absence of $K3$, the D4-branes and the O4-planes, this arrangement results in gauge group $SO(16) \times SO(16)$ for the type IA theory, resulting from the 16 D8-branes and one O8-plane at each end of the interval. As pointed out in refs.[18,19], this unique arrangement is suggestive: Away from the world volumes of the branes, *i.e.*, in the bulk of the x^5 interval, the model is locally the physics of the type IIA string. As we go to strong coupling in this model therefore, it is to be expected that a new dimension, x^{10} , opens up, and the model becomes an eleven dimensional theory. Each of the 16 D8-branes and O8-plane arrangements becomes a single object at strong coupling, supplying the two ends of the Hořava–Witten compactification of M-theory, becoming a sort of ‘nine-brane’ after unwrapping a leg in the x^{10} direction. Crucially, soliton states[20] which transform in the $(\mathbf{1}, \mathbf{128}) + (\mathbf{128}, \mathbf{1})$ (*i.e.*, the spinor of $SO(16) \times SO(16)$) become light in this limit, joining the $(\mathbf{1}, \mathbf{120}) + (\mathbf{120}, \mathbf{1})$ in filling out each $SO(16)$ into E_8 .

In fact, one can go further than this and see that *any other arrangement* of D8-branes and O8-planes will not produce an eleven dimensional theory. This issue is intimately related to the matter of when it is consistent to have D8-branes in type IIA theory⁹.

An isolated D8-brane in type IIA string theory is not consistent. As pointed out in refs.[19], the linear behaviour of the dilaton with the transverse distance is such that the string coupling diverges a *finite* distance away from the brane. In order to fix this state of affairs, it is necessary to introduce an object which acts as a sink for the dilaton field, cutting off the growth of the string coupling at some chosen reference finite value, λ_{IA}^c . By supersymmetry, the object with the correct charges to do this is the object with opposite R-R charge to the D8-brane, the O8-plane.

Once we have made a reference choice for the value λ_{IA}^c at which the type IIA string coupling should be cut off, we have fixed the distance that the O8-plane is from the D8-brane. Due to the -16 units of charge of an O8-plane, overall consistency requires there to be 16 D8-branes. On introduction of the O8-planes and D8-branes, we are no longer in the type IIA theory (globally) but the type IA theory. Between each D8-brane, the theory

⁹ I have benefited from conversations with R. Myers concerning the O8-plane matters discussed in the rest of this sub-subsection.

is a piece of type IIA string theory with a dilaton gradient. (At low energy, it is simply the massive type IIA supergravity of Romans[22,23].). We have constructed the physics of the neighbourhood of an endpoint of the type IA interval $\mathcal{I}$.

So the answer to the question “When is it consistent to have a D8-brane in type IIA theory?” has the answer “When it is in the type IA string theory”.

Now if we chose to increase the string coupling, and try to study what happens at strong coupling, the situation is interesting. In order to keep the theory (in the limit of infinite coupling) from developing isolated regions of infinite string coupling again, we preserve our chosen reference cutoff value λ_{IA}^c by simply moving the orientifolds closer to the D8-branes, as the coupling increases. We can do this indefinitely. However, there comes a point at which the interval upon which all of this physics is taking place is forced by this process to become so short (compared to $\sqrt{\alpha'}$) that we make stringy sense of it by simply T-dualizing along that interval to go to the type IB string where the distances in the dual direction are large compared to $\sqrt{\alpha'}$.

The resulting physics of the continued process of increasing the string coupling further until it is infinitely strong, is addressed succinctly by going to the dual theory to the type IB string — the $SO(32)$ heterotic string.

This process of increasing the string coupling in the type IA theory will always result — for any configuration of D8-branes and O8-planes except one— in the weakly coupled physics of the $SO(32)$ heterotic string in the limit, because the O8-planes and D8-branes will always have to be moved closer to one another until the physics is best addressed in terms of strong coupling type IB physics.

The exception to this situation is the unique case where the dilaton charges are canceled locally, as mentioned a few paragraphs hence. In that case, there is no need to move O8-planes (reducing the length of the interval) to cutoff regions of infinite coupling. The strong coupling physics of that configuration is related to that of the $E_8 \times E_8$ heterotic string, as discussed above.

2.3.2. Inclusion of $K3$, D4-Branes and O4-Planes

Returning to the $K3$ compactification, we see that the discussion of the last section has a counterpart here. In the compactified model, we have also to include D4-branes to cancel the charge of the O4-plane fixed points on $K3$.

In analogy with similar processes as for D-branes, one may think of the origin of the O4-planes as arising from wrapping the O8-plane¹⁰ on the $K3$, whose curvature is localized at its fixed points in this orbifold limit. $K3$ ’s non-zero value of $R \wedge R$ (supported at each fixed point) in the worldvolume action of the O8-plane endows each fixed point with O4-plane charge.

¹⁰ This way of thinking of O4-planes was suggested to me by R. Myers

This necessity to introduce D4-branes and O4-planes is entirely analogous to the compactification of M-theory on $K3 \times \mathcal{I}$, or the $E_8 \times E_8$ heterotic string on $K3$. The curvature of $K3$ requires the introduction of 24 instantons for consistency. In this type IA situation, the instantons on $K3$ are played by combinations of O4-planes and D4-branes. In the strong coupling limit, where we make contact with M-theory on $K3 \times \mathcal{I}$, we expect them to become five-brane like objects (they reveal a hidden leg wrapped around the new dimension x^{10} which opens up), destined to become E_8 instantons in the heterotic limit. This is the analogue of the way that the combination of 16 D8-branes and 1 O8-plane became the M-theory “nine-brane” boundary carrying the E_8 degrees of freedom. An isolated dynamical unit of four D4-branes would become an ordinary M5-brane, while inside the $K3$, a single O4-plane is related to half an M5-brane localized on a fixed point of $K3$.

Recalling that this particular orientifold model is expected to become the (12, 12) heterotic string compactification, we may deduce that it is precisely 16 D4-branes plus 16 O4-planes which together become 12 E_8 instantons: 1 D4 and 1 O4 is equivalent to $3/4$ an E_8 instanton. In particular, the O4-plane is equivalent to $1/2$ of an instanton, and the single D4-brane sitting in it in our special configuration is $1/4$ of an instanton. This is consistent with the T_5 -duality relation to the GP model and its relation to the $Spin(32)/\mathbb{Z}_2$ heterotic string compactification: There, four D5-branes make up one dynamical unit which is dual to a small heterotic instanton, a numerical relationship which should be invariant under T_5 -duality. Meanwhile, a single O5-plane is equivalent to a heterotic instanton, which is consistent with our type IA assignment above, and the fact that T_5 -duality turns two O4-planes into one O5-plane.

The relationship between the parameters of the M-theory compactification and those of type IA may be easily computed to be (see footnote 8):

$$\begin{aligned} L_M^{10} &= \lambda_{IA}^{\frac{2}{3}} \\ L_M^5 &= \lambda_{IA}^{-\frac{1}{3}} L_{IA}^5 \\ R_M &= \lambda_{IA}^{-\frac{1}{3}} R_{IA}, \end{aligned} \tag{2.5}$$

where $L_M^{10}/2\pi$, $L_M^5/2\pi$ and R_M are the radii of x^{10} direction, the x^5 direction and the $K3$, measured in M-theory length units. These relationships are simple consequences of dimensional reduction of the eleven dimensional supergravity Lagrangian on a circle to ten dimensions, and then performing a Weyl rescaling to match the resulting Lagrangian to the string frame Lagrangian of the massless sector of the relevant string theory[24].

Using the relations (2.4), we can compute the relationship between the M-theory param-

eters and the type IB parameters:

$$\begin{aligned}
L_M^{10} &= \frac{\lambda_{IB}^{\frac{2}{3}}}{(L_{IB}^5)^{\frac{2}{3}}} \\
L_M^5 &= \frac{1}{\lambda_{IB}^{\frac{1}{3}}(L_{IB}^5)^{\frac{2}{3}}} \\
R_M &= \frac{R_{IB}(L_{IB}^5)^{\frac{1}{3}}}{\lambda_{IB}^{\frac{1}{3}}}.
\end{aligned} \tag{2.6}$$

2.4. The $E_8 \times E_8$ and $Spin(32)/\mathbb{Z}_2$ Heterotic Strings on $K3$

At this stage we can contemplate moving back to six dimensions and recover the $E_8 \times E_8$ heterotic string compactified on $K3$. This is performed by starting with our M–theory configuration and shrinking the length of the interval by sending $L_M^5 \rightarrow 0$. (Shrinking $L_M^{10} \rightarrow 0$ would return us to the six dimensional theory of the type IA compactification.)

In the limit, we obtain the (12,12) heterotic compactification. This is guaranteed by fact that the in type IA orientifold model we chose an arrangement of branes which is completely symmetric between the ends of the interval. The tracing that we did of the instanton assignments from their role as $Spin(32)/\mathbb{Z}_2$ instantons in the type IB model, through the type IA model, and on to the M–theory configuration is therefore correct.

The relationship between the parameters of the heterotic model and those of the M–theory compactification are:

$$\begin{aligned}
L_M^5 &= \lambda_{HA}^{\frac{2}{3}} \\
L_M^{10} &= \lambda_{HA}^{-\frac{1}{3}} L_{HA}^{10} \\
R_M &= \lambda_{HA}^{-\frac{1}{3}} R_{HA},
\end{aligned} \tag{2.7}$$

where $L_{HA}^{10}/2\pi$, $L_{HA}^5/2\pi$ and R_{HA} are the radii of x^{10} direction, the x^5 direction and the $K3$, measured in $E_8 \times E_8$ heterotic string length units, while λ_{HA} denotes that heterotic string’s coupling¹¹.

Finally we arrive at the heterotic model with the assignment of instantons we expected from considerations[6] of $Spin(32)/\mathbb{Z}_2$ heterotic/type IB strong/weak coupling duality combined with T_{10} –duality between the two $K3$ compactified types of heterotic strings.

¹¹ For succinct subscripts it is simpler —and mindful of type I/heterotic duality— to use “HA” to denote the $E_8 \times E_8$ heterotic string and “HB” to denote the $Spin(32)/\mathbb{Z}_2$ heterotic string.

The T_{10} -duality gives the $Spin(32)/\mathbb{Z}_2$ heterotic string with parameters

$$\begin{aligned} L_{\text{HB}}^{10} &= \frac{1}{L_{\text{HA}}^{10}} \\ \lambda_{\text{HB}} &= \frac{\lambda_{\text{HA}}}{L_{\text{HA}}^{10}} \\ R_{\text{HB}} &= R_{\text{HA}}, \end{aligned} \tag{2.8}$$

where the notation should now be clear.

Combining this with (2.7) gives a relationship between the $Spin(32)/\mathbb{Z}_2$ heterotic parameters and those of M-theory:

$$\begin{aligned} L_{\text{M}}^5 &= \frac{\lambda_{\text{HB}}^{\frac{2}{3}}}{(L_{\text{HB}}^{10})^{\frac{2}{3}}} \\ L_{\text{M}}^{10} &= \frac{1}{\lambda_{\text{HB}}^{\frac{1}{3}} (L_{\text{HB}}^{10})^{\frac{2}{3}}} \\ R_{\text{M}} &= \frac{R_{\text{HB}} (R_{\text{IB}}^{10})^{\frac{1}{3}}}{\lambda_{\text{HB}}^{\frac{1}{3}}}. \end{aligned} \tag{2.9}$$

Comparing this to (2.6), if we did not know it already we could deduce[18] a strong/weak coupling duality between the type IB and the $Spin(32)/\mathbb{Z}_2$ heterotic string with the precise relationship[24,25,26]:

$$\begin{aligned} \lambda_{\text{IB}} &= \lambda_{\text{HB}}^{-1} \\ L_{\text{HB}}^5 &= L_{\text{HB}}^{10} \lambda_{\text{HB}}^{-\frac{1}{2}}, \end{aligned} \tag{2.10}$$

supplemented by the following relationship between the radii of the $K3$'s upon which they are compactified:

$$R_{\text{HB}} = R_{\text{IB}} \lambda_{\text{IB}}^{-\frac{1}{2}}. \tag{2.11}$$

For completeness, we also deduce a relationship between the type IA and $E_8 \times E_8$ parameters:

$$\begin{aligned} \lambda_{\text{IA}} &= \lambda_{\text{HA}}^{-\frac{1}{2}} (L_{\text{HA}}^{10})^{\frac{3}{2}} \\ L_{\text{IA}}^5 &= (L_{\text{HA}}^{10})^{\frac{1}{2}} \lambda_{\text{HA}}^{\frac{1}{2}}, \end{aligned} \tag{2.12}$$

with the $K3$'s radii related as follows:

$$R_{\text{IA}} = R_{\text{HA}} (L_{\text{HA}}^{10})^{\frac{1}{2}} \lambda_{\text{HA}}^{-\frac{1}{2}}. \tag{2.13}$$

2.5. Geometrical Picture of the Dualities

Taking into account all of the steps we have performed in constructing the chain of dualities, it should now be clear that the organizing geometry is that of a cylinder[18]. Essentially,

we compactify M–theory on the cylinder given by $S^1_{10} \times S^1_5 / \mathbb{Z}_2$, with length L_M^{10} for the ordinary circle and length L_M^5 for the line interval.

Shrinking L_M^{10} to zero sends us to the type IA string theory limit while shrinking L_M^5 sends us to the $E_8 \times E_8$ heterotic limit. Once in the string theory limit, T –duality along the surviving direction of the cylinder takes us to type IB string theory in case (x^5) , or $Spin(32)/\mathbb{Z}_2$ heterotic string theory in the other case (x^{10}) .

The strong/weak duality exchange of the type IB and $Spin(32)/\mathbb{Z}_2$ heterotic string is simply the exchanges of the values of L_M^5 and L_M^{10} , taking one from an infinitely thin line segment in one the first case to an infinitely short cylinder —the x^{10} circle— in the other.

2.6. Origin of Heterotic/Heterotic Duality

So far, we have not mentioned the extra special property this family of vacua have, the property which gives rise to heterotic/heterotic duality in six dimensions.

Its origin is clear once we realize that there is an alternative route from the type IB model to a heterotic model. Before going to type IA, we could perform the T –duality operation T_{6789} , which has the effect of defining a very similar type IB model, but on a $K3$ manifold obtained by inverting the radii of all of the circles in the defining T^4 . In fact, the model is invariant under this operation, as can be seen by examining the orientifold group[9].

Indeed, it is instructive to consider (separately) two interesting D–string probes in the six uncompactified directions, one a D1–brane, lying in the (x^0, x^4) directions, and another a D5′–brane wrapped about the $K3$ giving a string lying in the (x^0, x^5) direction.

type	x^0	x^1	x^2	x^3	x^4	x^5	x^6	x^7	x^8	x^9
D1	—	•	•	•	—	•	•	•	•	•
D5 (D1′)	—	•	•	•	•	—	—	—	—	—

Table 3. Orientation of D–brane probes in the orientifold model of Table 1 representing type IB string theory on $K3$, giving rise to two D–strings in six dimensions.

The operation T_{6789} defines type IB on the “dual” $K3$ which we call $\widetilde{K3}$, with radius $1/R_{IB}$. T_{6789} simply turns the D5–brane into a D1–brane and *vice-versa*, therefore exchanging our two types of D–string. Their tensions are easily computed[6] in terms of the type IB and the $Spin(32)/\mathbb{Z}_2$ heterotic parameters, using eqn.(2.11):

$$\begin{aligned} \mathcal{T}_{D1} &= \frac{1}{\lambda_{IB}} = \frac{R_{HB}^2}{R_{IB}^2} \\ \mathcal{T}_{D1'} &= \frac{R_{IB}^4}{\lambda_{IB}} = R_{HB}^2 R_{IB}^2, \end{aligned} \tag{2.14}$$

and are clearly exchanged under T_{6789} .

In purely $Spin(32)/\mathbb{Z}_2$ heterotic string paramters, their tensions are (respectively)[6]:

$$\begin{aligned}\mathcal{T}_{\text{electric}} &= R_{\text{HB}}^2 \lambda_{\text{HB}}^6 \\ \mathcal{T}_{\text{magnetic}} &= \frac{R_{\text{HB}}^2}{\lambda_{\text{HB}}^6},\end{aligned}\tag{2.15}$$

where $\lambda_{\text{HB}}^6 = R_{\text{IB}}^{-2}$ is the six dimensional heterotic string coupling. We see that the D1-brane becomes an electric “fundamental” heterotic string, being light at weak heterotic coupling. Meanwhile the D1’-brane becomes a magnetic “soliton” string, being heavy at weak coupling. Type IB’s T_{6789} -duality, which inverts R_{IB} , maps here to a strong/weak coupling duality.

Notice also that although we exchange two $K3$ ’s (with inversely related volumes) in the type IB model, the $K3$ which the heterotic string is compactified on remains the same, as can be seen by using eqn.(2.11), with the T_{6789} -duality relations

$$T_{6789} : \begin{cases} R_{\text{IB}} \rightarrow R_{\text{IB}}^{-1} \\ \lambda_{\text{IB}} \rightarrow \lambda_{\text{IB}}/R_{\text{IB}}^4 \end{cases}, \tag{2.16}$$

revealing that R_{HB} is invariant under the duality.

The orientations of the strings are trivially related to those in the first type IB model by a rotation by $\pi/2$, O_{45} , in the (x^4, x^5) plane.

type	x^0	x^1	x^2	x^3	x^4	x^5	x^6	x^7	x^8	x^9
D5 (D1’)	—	•	•	•	—	•	—	—	—	—
D1	—	•	•	•	•	—	•	•	•	•

Table 4. Orientation of D-brane probes of an orientifold representing type IB on $\widetilde{K3}$ (see Table 1), which is T_{6789} dual to the orientations of Table 3. These probes also give rise to two D-strings in six dimensions.

Instead of doing the operation T_{6789} we may proceed to our type IA model, using T_5 -duality. We thus obtain type IA on $K3 \times S_5^1/\mathbb{Z}_2$. Here our two types of brane probes become a D2-brane (with one leg stretched between the end of the x^5 interval) and a wrapped (on $K3$) D4-brane. (See Table 5)

type	x^0	x^1	x^2	x^3	x^4	x^5	x^6	x^7	x^8	x^9
D2	—	•	•	•	—	[—]	•	•	•	•
D4	—	•	•	•	•	•	—	—	—	—

Table 5. Two types of D-brane probe in the orientifold model of Table 2 realizing type IA on $K3 \times S_5^1/\mathbb{Z}_2$.

These two arrangements of branes (and the complete orientifold model) are dual to each other under T_{456789} -duality, which inverts the $K3$ and results in a model of type IA on $\widetilde{K3} \times S_4^1/\mathbb{Z}_2$. See Table 6. (A final $\pi/2$ rotation O_{45} would trivially restore the identical orientation.) This model is T_4 -dual to the type IB on $\widetilde{K3}$ model of Table 4.

type	x^0	x^1	x^2	x^3	x^4	x^5	x^6	x^7	x^8	x^9
D4	—	•	•	•	•	•	—	—	—	—
D2	—	•	•	•	[—]	—	•	•	•	•

Table 6. D-brane probe arrangements T_{456789} -dual to those in table 5. The background is an orientifold model of type IA on $\widetilde{K3} \times S_4^1/\mathbb{Z}_2$.

Upon moving to strong type IA coupling, we get two apparently distinct M-theory models, one on $K3 \times S_5^1/\mathbb{Z}_2$ and the other with $\widetilde{K3} \times S_4^1/\mathbb{Z}_2$. Our D2-brane probe becomes an M2-brane with one leg wrapped in the line interval $\mathcal{I}_5$ ($\mathcal{I}_4$), and transverse to the $K3$ (or $\widetilde{K3}$). The D4-brane becomes an M5-brane wrapped on the $K3$ (or $\widetilde{K3}$) and transverse to the $\mathcal{I}_5$ (or $\mathcal{I}_4$). See Tables 7 and 8.

type	x^0	x^1	x^2	x^3	x^4	x^5	x^6	x^7	x^8	x^9	x^{10}
M2	—	•	•	•	—	[—]	•	•	•	•	•
M5	—	•	•	•	•	•	—	—	—	—	—

Table 7. Two configurations of M-theory brane on $K3 \times \mathcal{I}_5$.

type	x^0	x^1	x^2	x^3	x^4	x^5	x^6	x^7	x^8	x^9	x^{10}
M5	—	•	•	•	•	•	—	—	—	—	—
M2	—	•	•	•	[—]	—	•	•	•	•	•

Table 8. *Two configurations of M-theory brane on $\widetilde{K3} \times \mathcal{I}_4$.*

From either one of these M-theory models, we may proceed to recover the heterotic string model by shrinking the line interval as discussed before. In both cases, the M2-brane is seen to become the light, electric, “fundamental” heterotic string (“F1-brane”) in the six dimensions $(x^0, x^1, x^2, x^3, x^{4(5)}, x^{10})$. Meanwhile, the M5-brane, wrapped on the $K3_M$ ($\widetilde{K3}_M$) become a wrapped (on $K3_H$) heterotic fivebrane soliton[27] (“F5-brane”) yielding a heavy solitonic string in the six dimensions. See Table 9.

type	x^0	x^1	x^2	x^3	x^4	x^6	x^7	x^8	x^9	x^{10}
F1	—	•	•	•	—	•	•	•	•	•
F5	—	•	•	•	•	—	—	—	—	—

Table 9. *The orientation of the two branes (an F1-brane and an F5-brane) which give rise to two dual $E_8 \times E_8$ heterotic strings in six dimensions. One is a light “fundamental” string, while the other is a heavy solitonic string.*

This heterotic model is compactified on $K3_H$, with six dimensional coupling $\lambda_H^6 \sim R_{IB}^{-2}$. Strong/weak coupling duality takes this to another model (with coupling $1/\lambda_H^6$) compactified on the same surface, with an exchange of the F1- and F5-branes. After a trivial relabeling of the x^5 coordinate to x^4 , we see that this is the model arising from reduction of the other M-theory configuration (table 10):

type	x^0	x^1	x^2	x^3	x^5	x^6	x^7	x^8	x^9	x^{10}
F5	—	•	•	•	•	—	—	—	—	—
F1	—	•	•	•	—	•	•	•	•	•

Table 10. *Branes which give dual $E_8 \times E_8$ heterotic strings in six dimensions.*

The two isomorphic tables above (9 and 10) are (by construction) respectively mapped

directly under the strong/weak coupling duality of type IB and heterotic strings¹² to the type IB models of tables 3 and 4, where x^5 (x^4) is exchanged with x^{10} .

So the T_{6789} -duality map between the two type IB models is pulled back to the strong/weak coupling¹³ heterotic/heterotic duality of Duff, Minasian and Witten[4], as pointed out in ref.[6]. Here, we have traced its action all the way through M-theory.

2.7. Heterotic/Heterotic Duality as Eleven Dimensional Electromagnetic Duality

Returning to the two M-theory models (tables 7 and 8) momentarily, it is natural to ask about the nature of the map between them. In the type I theories it was simply T -duality at work, while in the heterotic case it was six-dimensional electromagnetic strong/weak coupling duality. These must translate into something else in the full M-theory.

The answer is that as eleven dimensional models, the two models are electromagnetic duals of each other. It was suggested in ref.[4] that the operation of shrinking the interval $\mathcal{I}$ with the M2-brane wrapped on it was dual to another operation. Performing an eleven dimensional electromagnetic duality operation (with respect to the potential $A^{(3)}$) involves exchanging the sources, replacing the M2-brane with an M5-brane. Shrinking the M2-brane on $\mathcal{I}$ is then expected to be equivalent to wrapping the M5-brane on $K3$ and shrinking the volume of the $K3$. This conjecture was partially strengthened in ref.[4] by noting that precisely the (12,12) arrangement of instantons was needed to allow an M5-brane wrapped $K3$ to be shrunk completely.

However, there were geometric complications with making the construction completely precise. In order to argue that the result is again a heterotic string on $K3$, ref.[4] uses the intermediate seven dimensional duality between M-theory on $K3$ and the heterotic string on T^3 , and then reduces to six dimensions on the interval $\mathcal{I}$. This certainly gives a heterotic string in six dimensions, but gives no complete geometrical understanding of how the transverse $T^3 \times \mathcal{I}$ really becomes a $K3$ again, much less the *same* $K3$ on which the dual string is propagating.

The construction we have just performed in the previous subsection is a means of making the geometric proof complete. The key is that precisely for this situation, the details of the electromagnetic duality between the required configurations as eleven dimensional models are *completely* implied by the T_{456789} duality between the two type IA models, with their accompanying brane configurations depicted in tables 6 and 7.

The electromagnetic duality should be expected to work as follows: The M2-brane couples

¹² Strictly, as discussed earlier, we must also perform a T_{10} duality operation to move from $E_8 \times E_8$ to $Spin(32)/\mathbb{Z}_2$ heterotic.

¹³ It could not have mapped to a T -duality, as there is no such symmetry between the F1-brane and the F5-brane.

to the potential $A^{(3)}$ electrically and carries the basic unit of charge, while the M5-brane couples to it magnetically, being the dual object to the M2-brane in eleven dimensions.

In otherwise flat eleven dimensions with branes of infinite extent, the operation of replacing $A^{(3)}$'s field strength $F^{(4)}$ by its dual $F^{(7)}$, while replacing M2-branes by M5-branes (and *vice-versa*), would be a symmetry of the theory.

Here however, things are slightly more complicated. We have M-theory on $K3_M \times \mathcal{I}_5$, with sizes R_M and L_M^5 , respectively. The M2-brane is an open M2-brane, as it has boundaries on the ends of the interval $\mathcal{I}_5$. Part of its volume is finite, with a scale set by L_M^5 . If electromagnetic duality is going to work, we must conclude that the world volume of the dual M5-brane must also have a finite part. The scale of that volume must be a monotonically increasing function of L_M^5 , if we are to recover the uncompactified situation of the previous paragraph in the limit.

This finite part of the M5-brane world volume is wrapped on a *different* $K3$ from the one in the starting configuration, which we can call $\widetilde{K3}_M$. The radius $\widetilde{R}_M$ must be an increasing function of L_M^5 . (It has to be a different $K3$, as the other one must stay fixed while we reduce the M2-brane-wrapped interval $\mathcal{I}_5$.)

Similarly, had we started with an M5-brane wrapped on the $K3_M$, with radius R_M we must conclude that the dual configuration is an M2-brane wrapped on a line interval we might call $\mathcal{I}_4$, with length L_M^4 an increasing function of R_M , and transverse to a *different* $K3$ called $\widetilde{K3}_M$.

Precisely this situation is what we have in tables 7 and 8. The well-established T_{456789} -duality between these two configurations tells us precisely how the eleven dimensional duality must work. The parameters of one M-theory configuration are given by the equations (2.5) relating them to the parameters of the type IA configuration in table 5. Meanwhile, upon using the relations of T_{456789} duality between table 5 and table 6 (together with $L_M^4 = \lambda_{IA}^{-1/3} L_{IA}^4$):

$$\begin{aligned} R_{IA} &\leftrightarrow \frac{1}{R_{IA}} \\ L_{IA}^{5(4)} &\leftrightarrow \frac{1}{L_{IA}^{5(4)}} \\ \lambda_{IA} &\leftrightarrow \frac{\lambda_{IA}}{L_{IA}^4 L_{IA}^5 (R_{IA})^4} \end{aligned} \tag{2.17}$$

we can deduce the parameters of the dual M-theory configuration to be related thus:

$$\begin{aligned} \widetilde{R}_M &= (L_M^5 L_M^4 R_M)^{\frac{1}{3}} \\ \widetilde{L}_M^{4(5)} &= \left[(R_M)^4 \frac{L_M^{5(4)}}{(L_M^{4(5)})^2} \right]^{\frac{1}{3}}. \end{aligned} \tag{2.18}$$

So far these relations are apparently in line with our expectations above that shrinking $\mathcal{I}_5$ is dual to shrinking $\widetilde{K3}_M$, and also that shrinking the dual interval $\mathcal{I}_4$ is equivalent to

shrinking $K3_M$. However, we must be careful. At face value, they also seem to suggest that shrinking $\mathcal{I}_5$ results in the shrinking of $\mathcal{I}_4$, which certainly does not seem correct.

To set things aright, it is illuminating to factor out of the expressions the quantity $V_M \equiv L_M^4 L_M^5 (R_M)^4$, the volume of the compact part $K3_M \times \mathcal{I}_5 \times S_4^1$ of the M–theory compactification, whose image in type IA we T_{456789} –dualize. Doing this, we obtain:

$$\begin{aligned} \tilde{R}_M &= \frac{V_M^{\frac{1}{3}}}{R_M} \\ \tilde{L}_M^{4(5)} &= \frac{V_M^{\frac{1}{3}}}{L_M^{4(5)}} \\ \text{and also } \tilde{L}_M^{10} &= \frac{V_M^{-\frac{2}{3}}}{L_M^{10}}. \end{aligned} \tag{2.19}$$

These relations look like an M–theory analogue of type IA’s T_{456789} –duality, with a volume factor entering in an unusual way (*c.f.* eqn.(2.17)). Notice however that the dual volume is the same as the original volume:

$$\tilde{V}_M \equiv \tilde{L}_M^4 L_M^5 (\tilde{R}_M)^4 = \frac{V_M^{\frac{1}{3}}}{L_M^4} \frac{V_M^{\frac{1}{3}}}{L_M^5} \frac{V_M^{\frac{4}{3}}}{(R_M)^4} = V_M, \tag{2.20}$$

and so the apparent length inversions in eqn.(2.19) are internal rearrangements within the “dualized” surface, performed *at fixed volume*. This is closer to what we should expect from the electromagnetic duality, and less like a stringy T –duality.

We therefore interpret equation (2.19) as follows. First, we will keep the volume V_M fixed while shrinking any part of $K3_M \times \mathcal{I} \times S^1$. Then, shrinking $\mathcal{I}_5$, we must rescale the radius R_M of the $K3_M$ to keep the volume fixed. In the dual M–theory configuration, this is equivalent to (according to equation (2.19)) shrinking $\tilde{R}_M$, the radius of the dual $\tilde{K}3_M$. The same reasoning may be used to relate the shrinking of the dual interval $\mathcal{I}_4$ to the shrinking of $K3_M$ in the original geometry.

So we have sharpened the statement of Duff, Minasian and Witten concerning the eleven dimensional origin of heterotic/heterotic duality: Wrapping an M2–brane on the interval $\mathcal{I}_5$ and shrinking the interval to zero size to obtain a heterotic string compactified on $K3_H$ to six dimensions, is equivalent (by electromagnetic duality) to wrapping an M5–brane on $\tilde{K}3_M$ and shrinking the volume of the $\tilde{K}3_M$ to zero to get a dual heterotic string compactified on the *same* $K3_H$ to six dimensions.

In summary, our main tool to study the details of the eleven dimensional duality was to realize it as the image of T–duality between two type IA configurations. Those configurations extend to two apparently different M–theory compactifications under the strong coupling map, and the duality map between the two M–theory configurations follows from

the existence of the T-duality map composed with the (unique) strong/weak coupling map between of type IA and M-theory on a circle.

In this way, we also see that there is no mystery of how we recover $K3_H$ for the resulting heterotic string in the case of the $K3_M$ -shrink-wrapping of the M5-brane, *and* we see that we recover the correct $K3_H$ as well.

DMW's result that there is no obstruction to shrinking the dual M5-brane on a $K3$ when there is a (12,12) instanton compactification directly translates into the statement that T_{456789} -duality relates two type IA configurations which can be taken to M-theory at strong coupling, and thus in turn becomes eleven dimensional electromagnetic duality in the limit.

3. Probing with D-Branes

Let us now return to the study of the full details of the type I orientifolds model defined in tables 1 and 2. At the bottom of table 2 is shown the orientation of a D0-brane probe which we might introduce into the model.

This is the starting point for the derivations of the various models which we use to motivate parts of a matrix theory representation of the $K3_M \times \mathcal{I}$ compactification of M-theory. To derive a matrix theory we will use not just one D0-brane, but N of them. Following Banks, Fischler, Shenker and Susskind[12], and in the notation of the previous section, the large N limit will yield a matrix theory representation of M-theory on $K3_M \times \mathcal{I}_5$ in the infinite momentum frame, where the infinite boost is in the x^{10} direction.

The logic of the procedures will be as follows: Considering first only the local geometry for a moment, we see that we can derive (as the effective D0-brane world-line theory) a model of 0+1 dimensional quantum mechanics which encapsulates the physics of the probe moving around in the background of the D8- and D4-branes, the O8- and 04-planes, and the $K3$. Given the strong coupling discussions in section 2.3.2, in the matrix model (and large N) limit, such a 0+1 dimensional model should capture aspects of the M-theory physics of small E_8 instantons on an ALE space, near an end of the interval $\mathcal{I}_5$.

In moving on to consider the case where the compactness of the $\mathcal{I}_5$ comes into play, we should expect that we must take into account light strings interconnecting the N D0-branes after first wrapping this interval when it is small. Operationally, (after a Fourier transform in the matrix theory language) this gives us a 1+1 dimensional model, derivable as the physics of the T_5 -dual situation with N S_5^1 -wrapped D1-brane probes, oriented as at the bottom of table 1.

This interlaces nicely with the fact that a D1-brane probe in type IB string theory is the heterotic string[25]: In the $L_5 \rightarrow 0$ limit, the matrix theory is an uncompactified 1+1 dimensional model. This 1+1 dimensional matrix theory is the matrix theory of the

heterotic string on a local piece of $K3$, *i.e.*, near an ALE space, with (part of) an E_8 instanton on it. In the strong coupling limit, which is the IR limit of the model, we expect to find a conformal field theory of this string, along the lines of Dijkgraaf, Verlinde and Verlinde[28] for the recovery of ten dimensional type IIA strings from matrix theory¹⁴.

Furthermore, when we consider the next feature —that the $K3$ is also small and compact— we must take into account light strings coming from winding the $K3$. Again, we see that this is morally equivalent to taking the T_{6789} -dual of the D1-branes defining the 1+1 dimensional model and working with the world-volume theory of the dual probe, wrapped on a dual surface. In this way we see that we are working with the world volume theory of N D5'-branes shown at the bottom of table 1, and we therefore have (at least part of) a definition in terms of a particular 5+1 dimensional quantum theory compactified on the surface $\widetilde{K3} \times S_5^1$.

Notice that heterotic/heterotic duality is already manifest in this matrix theory definition. Recall from section 2 that this duality is generated by T_{6789} -duality, which in this context exchanges the N D1-branes with the N D5'-branes.

As this theory is related in this way to the world volume of a collection of N D5'-branes in type IB string theory, we suspect that it is related to the $(0,1)$ theory with $Spin(32)/\mathbb{Z}_2$ global symmetry[30]. We will thus be led to examine the details of that expectation (in section 4), and furthermore to consider what it means to compactify such a theory on $\widetilde{K3} \times S_5^1$. Much later, we will speculate that moving away from the special DMW/GP point, we should consider the matrix theory definition of M-theory on $K3 \times \mathcal{I}_5$ to be given by the $(0,1)$ $E_8 \times E_8$ theory compactified on the dual given by $\widetilde{K3} \times S_5^1$.

3.1. Probing with D1-Branes: The Model.

As a starting point, we study the field content of the world-volume theory of the N D1-brane probes in the type IB model. Returning to the type IA case with N D0-branes will be easily performed by (essentially) dimensional reduction.

We will refer to table 1 for the orientations of the GP model's branes and the $K3$ (top part) and the N D1-brane probes (bottom).

3.1.1 Supersymmetry

The D-brane orientations of the model (with D1-branes) breaks the Lorentz group up as follows:

$$SO(1,9) \supset SO(1,1)^{05} \times SO(4)^{1234} \times SO(4)^{6789}, \quad (3.1)$$

¹⁴ It may be interesting to see also the comments at the end of ref.[29], for some anticipation of this way of realizing duality.

where the superscripts denote the sub-spacetimes in which the surviving factors act. Following refs.[31,32], we may label the worldsheet fields according to how they transform under the covering group (which acts as an R-symmetry of our final 1+1 dimensional model):

$$G = [SU(2)' \times \widetilde{SU(2)}']_{1234} \times [SU(2)_R \times SU(2)_L]_{6789}, \quad (3.2)$$

with doublet indices $(A', \tilde{A}', A, Y)$, respectively.

We will place the $K3$ into the mix by embedding the reflection symmetry R_{6789} , into the $SO(4)_{6789}$.

Accordingly, the supercharges decompose under (3.1) first (due to the D-strings) as

$$\mathbf{16} = \mathbf{8}_+ + \mathbf{8}_- \quad (3.3)$$

where $\pm$ subscripts denote a chirality with respect to $SO(1,1)$, and furthermore (due to the D5-branes and the $K3$) each $\mathbf{8}$ decomposes into a pair of $\mathbf{4}$'s of the $SO(4)$'s. As we will see in due course, the various projections will pick out one of these spinors to carry the supersymmetry on the world sheet, giving a 1+1 dimensional system with $(0,4)$ supersymmetry, as we might expect from traditional heterotic string (on $K3$) considerations, keeping in the back of our minds that this is of course dual to that very system¹⁵.

The spectrum of massless fields in the model will produce a family of fields on the world-volume, which has coordinates (x^0, x^5) . The supersymmetry algebra is of the form:

$$\{Q^{AA'}, Q^{BB'}\} = \epsilon^{AB} \epsilon^{A'B'} P_-, \quad (3.4)$$

where $P_- \equiv -i\partial/\partial\sigma^-$, where $\sigma^\pm = (x^0 \pm x^5)/2$. Here, ϵ^{AB} and $\epsilon^{A'B'}$ are the antisymmetric tensors of $SU(2)$ and $SU(2)'$, respectively, A, B, A' and B' being doublet indices.

3.1.2 The 1-1 Strings

As there are 8 Dirichlet-Dirichlet (DD) directions, the Neveu-Schwarz (NS) sector has zero point energy $-1/2$. The massless excitations form vectors and scalars in 2D, and are created as follows. For the vectors, the Neumann-Neumann (NN) directions give excitations:

$$A^{0,5}(x^0, x^5) : \quad \lambda_V \psi_{-\frac{1}{2}}^{0,5} |0\rangle, \quad (3.5)$$

where λ_V is an $2N \times 2N$ Chan-Paton matrix, which satisfies (at a fixed point of R_{6789}):

$$\begin{aligned} \Omega : \quad \lambda_V &\rightarrow -\gamma_\Omega \lambda_V^T \gamma_\Omega^{-1} \\ R_{6789} : \quad \lambda_V &\rightarrow \gamma_R \lambda_V \gamma_R^{-1}, \end{aligned} \quad (3.6)$$

¹⁵ Notice that this is the same amount of supersymmetry as possessed by a system of D9- D5- and D1-branes[33]. The $K3$ does not break any more supersymmetry than the D5-branes already do[9].

where the $\gamma_{\Omega(R)}$ matrices are $2N \times 2N$ matrices chosen to represent the action of Ω and R_{6789} on λ_V . We will use the same type of basis as ref.[9]. The solution of the constraint equations (3.6) results in λ being generators of the group $U(N)$.

Away from a fixed point, the second constraint in eqn.(3.6) does not apply, and R_{6789} merely relates a D-brane to its image. Instead, only the orientation projection acts, giving gauge group $SO(N)$. This is consistent with the fact that Ω^2 has the same sign for D1-branes as for D9-branes[9].

The excitations coming from the 8 DD directions split into two parts. There are fields associated with excitations in the $x^{1,2,3,4}$ directions, and fields associated with the $x^{6,7,8,9}$ directions:

$$\begin{aligned}\phi(x^0, x^5) : \quad & \lambda_\phi \psi_{-\frac{1}{2}}^{1,2,3,4} |0\rangle > \\ \psi(x^0, x^5) : \quad & \lambda_\psi \psi_{-\frac{1}{2}}^{6,7,8,9} |0\rangle >,\end{aligned}\tag{3.7}$$

where $\lambda_{\phi(\psi)}$ are $N \times N$ Chan-Paton matrices satisfying (at a fixed point):

$$\begin{aligned}\Omega : \quad & \begin{cases} \lambda_\phi \rightarrow -\gamma_\Omega \lambda_\phi^T \gamma_\Omega^{-1} \\ \lambda_\psi \rightarrow \gamma_\Omega \lambda_\psi^T \gamma_\Omega^{-1} \end{cases}, \\ R_{6789} : \quad & \begin{cases} \lambda_\phi \rightarrow \gamma_R \lambda_\phi \gamma_R^{-1} \\ \lambda_\psi \rightarrow -\gamma_R \lambda_\psi \gamma_R^{-1} \end{cases}.\end{aligned}\tag{3.8}$$

(Once again, away from a fixed point, R_{6789} places no constraint, and simply reflects.)

Now since λ_ϕ satisfies the same constraints as λ_V , we see that the ϕ fields are a family of four-component scalars transforming in the adjoint of $U(N)$, (or of $SO(N)$ away from a fixed point), which we shall denote as $b^{A'\tilde{A}'}$, to make contact with refs.[31,32]. Meanwhile, the ψ fields are a complex doublet of hypermultiplet fields transforming in the antisymmetric representation of the gauge group $U(N)$ or of $SO(N)$. We denote them $b^{AY}(x^0, x^5)$, in accordance with refs.[31,32].

The fermionic states ξ from the Ramond (R) sector (with zero point energy 0, by definition) are built on the vacua formed by the zero modes ψ_0^i , $i=0, \dots, 9$. This gives the initial **16** of the left hand side of eqn. (3.3). The GSO projection acts on the vacuum in this sector as:

$$(-1)^F = \Gamma^0 \Gamma^1 \dots \Gamma^9,\tag{3.9}$$

while as Ω acts as -1 on NN strings (*i.e.*, in the (x^0, x^5) directions), it is:

$$\Omega = \Gamma^1 \Gamma^2 \Gamma^3 \Gamma^4 \Gamma^6 \Gamma^7 \Gamma^8 \Gamma^9,\tag{3.10}$$

and as R_{6789} is -1 in the $x^{6,7,8,9}$ directions it is:

$$R_{6789} = -\Gamma^6 \Gamma^7 \Gamma^8 \Gamma^9.\tag{3.11}$$

So we have $(-1)^F \xi = \xi$ from the GSO projection, and with Ω , it simply correlates world sheet chirality with spacetime chirality: $\Gamma^0 \Gamma^5 \xi_{\pm} = \pm \xi_{\pm}$, where ξ_- is in the $\mathbf{8}_c$ of $SO(8)$ and ξ_+ is in the $\mathbf{8}_s$.

Now ξ_- is further decomposed by R_{6789} into ξ_-^1 and ξ_-^2 , where superscripts 1 and 2 denote the decomposition into the (1234) sector and the (6789) sector, respectively. So we have that the four component fermion ξ_-^1 (hereafter called $\psi_-^{A\tilde{A}'}$) is the right-moving superpartner of the four component scalar field $b^{A'\tilde{A}'}$, while ξ_-^2 (called $\psi_-^{A'Y}$) is the right-moving superpartner of b^{AY} . The supersymmetry transformations are:

$$\begin{aligned}\delta b^{A'\tilde{A}'} &= i\epsilon_{AB}\eta_+^{A'A}\psi_-^{B\tilde{A}'} \\ \delta b^{AY} &= i\epsilon_{A'B'}\eta_+^{AA'}\psi_-^{B'Y}.\end{aligned}\tag{3.12}$$

The ψ carry the same $U(N)$ (or $SO(N)$) charges as their bosonic superpartners in order to survive the R_{6789} and Ω projections, thus ensuring that gauge symmetry respects supersymmetry.

Similarly, the field ξ_+ is decomposed under (3.1) into ξ_+^1 (which we'll call $\psi_+^{A\tilde{A}'}$) and ξ_+^2 (called $\psi_+^{A'Y}$). Formally, these fields are left-moving “superpartners” of the gauge field A^μ . We have only (0, 4) supersymmetry, and so no *linearly* realized left-moving supersymmetry transformations, but there are non-linear ones. As A^μ is not dynamical in two dimensions, the detailed form of these transformations will not concern us here¹⁶.

3.1.3 The 1–9 Strings

There are 8 Dirichlet–Neumann (DN) coordinates, giving ground state energy 1/2, and so there are no massless states arising in the NS sector. The R sector excitations come from the NN (x^0, x^5) system giving just two ground states. In this sector, the GSO projection is simply $(-1)^F = \Gamma^0 \Gamma^5$, which picks the left-moving field[25].

Assuming (for now) that we are in the configuration where we have maximal gauge group $U(16)_9$ from the D9–branes, we have left-moving fermions λ_+ in the $(\mathbf{N}, \mathbf{16})$, the first part of the current algebra fermions of the heterotic string. We denote a component by λ_+^M , where M is a D9–brane index, whenever we need to explicitly show the index structure under the global symmetry of the D9–brane gauge group.

(We shall see a little later that we have not completely determined the 1–9 spectrum. There is another sector which arises which will play a crucial role in the proceedings.)

¹⁶ In the case of a single D1–brane, A^μ and ξ_+ would be projected out by acting with Ω , leaving us with just the right-moving states ξ_- as superpartners of the b -fields, as in standard heterotic lore[25]. In this case, we have Chan Paton factors λ_V , on which a representation of Ω can act to supply us with a correctly transforming surviving state.

3.1.4 The 1–5 Strings

There are four DN coordinates, and four DD coordinates giving the NS sector a zero point energy of 0, with excitations coming from integer modes in the 1234 directions, giving a four component boson. This is sector 1, in the notation above. The R sector also has zero point energy of zero, with excitations coming from the 6789 directions, giving a four component fermion χ . This is sector 2, as above.

The GSO projections in either sector are:

$$\begin{aligned} (-1)_1^F &= \Gamma^0 \Gamma^1 \Gamma^2 \Gamma^3 \Gamma^4 \Gamma^5 \\ (-1)_2^F &= \Gamma^0 \Gamma^5 \Gamma^6 \Gamma^7 \Gamma^8 \Gamma^9, \end{aligned} \quad (3.13)$$

which, upon application, reduce us to two bosonic states $\phi^{A'}$ in sector 2, and decomposes the spinor χ in sector 1 into left and right moving two component spinors, χ_-^A and χ_+^Y , respectively. We see that χ_-^A is the right-moving superpartner of $\phi^{A'}$.

Assuming also that we have chosen the configuration where all of the D5-branes are coincident at one fixed point, we have therefore a supermultiplet in the $(\mathbf{N}, \mathbf{16})$, with components $(\phi^{A'm}, \chi_-^{Am})$:

$$\delta\phi^{A'm} = i\epsilon_{AB}\eta_+^{A'A}\chi_-^{Bm}. \quad (3.14)$$

Here, m is a D5-brane group theory index.

Also, (with components χ_+^{Ym}), χ_+^Y transforms in the $(\mathbf{N}, \mathbf{16})$. They are the rest of the current algebra fermions of the heterotic string, which from this point of view carry any non-perturbative global symmetry arising due to the D5-branes' behaviour as small instantons.

3.1.5 The 9–9, 5–5 and 9–5 Strings

Crucial to the whole discussion is that fact that these fields will generically appear as couplings in the 1+1 dimensional theory.

From the 5–5 sector there are four-component scalar couplings, (descendants of the $x^{6,7,8,9}$ hypermultiplets) transforming in the $\mathbf{120} + \overline{\mathbf{120}}$, which we call X_{mn}^{AY} , matching the notation of ref.[32].

There are similar fields in the 9–9 sector and we denote these couplings Y_{MN}^{AY} .

Meanwhile, the 9–5 sector produces a $(\mathbf{16}, \mathbf{16})$, denoted h_M^{Am} , with m and M showing off its choices in D5- and D9-brane group theory.

Now that we have named these fields, we can display the supersymmetry transformation relating them to the left moving fields:

$$\begin{aligned} \delta\lambda_+^M &= \eta_+^{AA'} C_{AA'}^M \\ \delta\chi_+^{Ym} &= \eta_+^{AA'} C_{AA'}^{Ym}, \end{aligned} \quad (3.15)$$

where:

$$\begin{aligned} C_{AA'}^M &= h_A^{Mm} \phi_{A'm} \\ C_{AA'}^{Ym} &= \phi_{A'}^n (X_{An}^{Ym} - b_A^Y \delta_n^m), \end{aligned} \quad (3.16)$$

These precise transformations allow us to write the non-trivial part of the $(0, 4)$ supersymmetric 1+1 dimensional Lagrangian containing the Yukawa couplings and the potential:

$$\begin{aligned} \mathcal{L}_{\text{tot}} = \mathcal{L}_{\text{kinetic}} - \frac{i}{4} \int d^2\sigma & \left[\lambda_+^M \left(\epsilon^{BD} \frac{\partial C_{BB'}^M}{\partial b^{DY}} \psi_-^{B'Y} + \epsilon^{B'D'} \frac{\partial C_{BB'}^M}{\partial \phi^{D'm}} \chi_-^{Bm} \right) \right. \\ & + \chi_+^{Ym} \left(\epsilon^{BD} \frac{\partial C_{BB'}^{Ym}}{\partial b^{DY}} \psi_-^{B'Y} + \epsilon^{B'D'} \frac{\partial C_{BB'}^{Ym}}{\partial \phi^{D'm}} \chi_-^{Bm} \right) \\ & \left. + \frac{1}{2} \epsilon^{AB} \epsilon^{A'B'} (C_{AA'}^M C_{BB'}^M + C_{AA'}^{Ym} C_{BB'}^{Ym}) \right]. \end{aligned} \quad (3.17)$$

This was derived in ref.[31] as the most general $(0, 4)$ supersymmetric Lagrangian with these types of multiplets, providing that the C satisfy the condition:

$$C_{AA'}^M C_{BB'}^M + C_{AA'}^{Ym} C_{BB'}^{Ym} + C_{BA'}^M C_{AB'}^M + C_{BA'}^{Ym} C_{AB'}^{Ym} = 0, \quad (3.18)$$

which they do[32]. $\mathcal{L}_{\text{kinetic}}$ contains the usual kinetic terms for all of the fields, and the required terms which complete them into gauge invariant terms.

It should be of some concern that the 9–9 hypermultiplet fields Y_{MN}^{AY} have not entered into the story in a non-trivial way. We see their role in what follows:

3.1.6 Two Puzzles, and Their Solution

Running a little ahead of the story for a minute, let us anticipate what this model should do for us thus far. We have written down the physics of the (multi-) D1-brane probe (slightly generalizing ref.[33,32]), and we expect (following refs.[32,31,34]) that the conditions (3.18) will restrict us to the moduli space of vacua of the 1+1 dimensional theory which is isomorphic to the moduli space of instantons. In other words, equations (3.18) are the ADHM data, and finishing the search for the gauge inequivalent vacua by setting the potential to zero and gauge fixing will compute a hyperKähler quotient for us. This makes contact with the fact[35] that the D5-branes¹⁷ are instantons in the D9-brane gauge fields, dual to heterotic instantons.

This anticipation is correct. However, our studies in section 2 should suggest that something is not quite right. Indeed, the D5-branes are correctly to be seen as instantons in the model, but they are not the only objects playing that role. As there are only 8 full instantons to be found in the D5-brane sector, the other 16 needed for a consistent $K3$

¹⁷ More precisely, a quartet of them[9]. See also section 2 in this paper.

compactification are to be found elsewhere: There is a single instanton living at the core of each fixed point of R_{6789} .

So even in the complete absence of D5-branes, it must be that the D1-brane probes should be able to detect instantons, by finding them in the moduli space of their world-sheet sigma model. In the model that we have above (eqn.(3.17)), if we crudely mimic the absence of D5-branes by setting all of the 1-5, 5-5 and 9-5 fields to zero¹⁸, we are left with a trivial model with not enough structure to give us —after the hyperKähler quotient— anything other than either the trivial flat space (over-endowed with a hyperKähler structure) when we are away from a fixed point, or the (resolved; see the next sub-subsection) ALE space.

So somehow things need to be modified in order that the D1-branes see an instanton hiding in the ALE space. This is our first puzzle.

This is not unrelated to the apparent absence of a role for the 9-9 fields Y_{MN}^{AY} . If they are to be non-trivially involved, they must somehow enter the lagrangian in a (0, 4) respecting way. The only way to do so is to enter in equations of the form (3.16). However, an examination of the index structure shows that there is nothing for Y_{MN}^{AY} to couple to in order to enter consistently.

Persevering, an examination of the λ_+^M supersymmetry transformations (3.15) which it must enter finds the source of the problem: As it stands the supersymmetry transformations are of the form 1-9=1-5·5-9 or 1-5=1-5·5-5, showing the fusion of D1-, D5-, and D9-index structure. There is no way to form such an equation with this structure using Y_{MN}^{AY} and the fields we already have: We need a 1-9 field with indices A', M . This constitutes our second puzzle.

If both problems find their solution in a common place it must be that something happens at the fixed points of R_{6789} . This is precisely at the heart of the first problem, as we know that away from the fixed points, the system should be identical to that which has gone before[32,31] for the 1-5-9 system in flat space.


Turning therefore back to our computation of the spectrum of the 1-9 sector we see that there is indeed something extra at a fixed point. For a D1-brane sitting at a fixed point, there is a wholly new sector of massless strings arising in the 1-9 sector. These new strings are special (*c.f.*, section 3.1.3) in that they have their endpoints fixed in the $x^{6,7,8,9}$ directions.

As the literature does not seem to elaborate on such strings, let us be explicit in showing where they come from: It is analogous to the case of the extra massless 1-1 strings which arise when a D1-brane alights upon a fixed point. First consider, in the covering space of the orbifold, the D1-brane at some position $\mathbf{X}=\mathbf{X}_0$ away from the fixed point which is (say) at $\mathbf{X}=0$. The massless 1-9 strings arise entirely from strings beginning on the

¹⁸ To be less crude, we would actually just send the masses of the 1-5 fields to infinity, telling us that the D5-branes are infinitely far away.

D1-brane and ending on the nearby D9-branes (which are of course everywhere) at $\mathbf{X}_0$. These strings will also contribute to the spectrum when the D1-brane is at the fixed point, and are considered in section 3.1.3.

There are however, 1–9 strings which stretch from the D1-brane to the mirror image position on the D9-branes, $-\mathbf{X}_0$ (near the mirror partner of the D1-brane, which is not relevant here). These are stretched, and do not contribute to the massless spectrum. On the orbifolded space, we must think of these strings as stretching from the D1-brane at $\mathbf{X}_0$ to the fixed point $\mathbf{X}=0$, and back to the vicinity of the D1-brane. Although it is back where it started, due to the reflection, it is stretched to length $2\mathbf{X}_0$, and looped through the fixed point. The figure shows both points of view schematically (we have let the strings touch the D9-branes slightly away from the D1-brane, for clarity):



As the D1-brane approaches the fixed point, these looped strings become of zero length and therefore contribute to the massless spectrum *in addition to the ordinary 1–9 strings*. The crucial difference between these “twisted 1–9 strings” and the garden variety is that they are trapped at the fixed point *i.e.*, in the $x^{6,7,8,9}$ directions. So instead of 8 DN coordinates, as computed previously in section 3.1.3, giving a zero point energy of $1/2$, there are 4 DN coordinates ($x^{1,2,3,4}$) and 4 DD coordinates ($x^{6,7,8,9}$), shifting the zero point energy to $-1/2$. Now we can excite massless states in both the NS and R sector, just as computed for the 1–5 sector previously¹⁹.

In this way, we get three new fields for the 1–9 sector when we are at a fixed point: the

¹⁹ Indeed, it is as if we have introduced a whole new type of five-brane which is trapped at the fixed point. We can think of this as an explicit orbifold realization of the phenomenon of wrapping a D9-brane on $K3$ to get a five-brane. This new five-brane is different from ordinary D5-branes, though. It couples to the closed string twist fields, which is another reason why it is stuck at the fixed points. The cylinder and Möbius strip diagrams of refs.[9.16] which have twisted sector fields propagating in the closed string channel, may be thought of as having these five-branes attached to their boundaries.

supermultiplet $(\rho^{A'M}, \zeta_-^{AM})$, with supersymmetry transformations:

$$\delta\rho^{A'M} = i\epsilon_{AB}\eta_+^{A'A}\zeta_-^{BM}, \quad (3.19)$$

and the left mover ζ_+^{YM} . (M is a D9-brane index.)

This is precisely what we need to solve our problem. At a fixed point, we have in addition to the term in the Lagrangian involving λ_+^M , we have a term

$$\zeta_+^{YM} \left(\epsilon^{BD} \frac{\partial C_{BB'}^{YM}}{\partial b^{D'Y}} \psi_-^{B'Y} + \epsilon^{B'D'} \frac{\partial C_{BB'}^{YM}}{\partial \phi^{D'M}} \zeta_-^{BM} \right), \quad (3.20)$$

where

$$C_{AA'}^{YM} = \rho_{A'}^N (Y_{AN}^{YM} - b_A^Y \delta_N^M), \quad (3.21)$$

and

$$\delta\zeta_+^{YM} = \eta_+^{AA'} C_{AA'}^{YM}, \quad (3.22)$$

which has a 1-9=1-9-9-9 structure matching the 1-5=1-5-5-5 structure of the second equation in (3.15). We complete the modification of our Lagrangian by adding obvious terms involving $C_{AA'}^{YM}$ to eqn.(3.18).

This modification has a number of virtues, in solving our problem:

- (i) It is the unique way that we can modify the 1-9 sector of the lagrangian and still keep the supersymmetry structure.
- (ii) We see how the fixed points break D1-D5 translational symmetry *via* the inclusion of the couplings (3.21). With the couplings (3.16) alone, there is translational symmetry which should only be appropriate away from the fixed points.
- (iii) We see now that even in the absence of all D5-brane fields, the D9-branes supply couplings which satisfy the same form of (ADHM) specification (3.18) that the D5-brane fields did. Therefore, the D1-branes will correctly see instantons in the D9-branes, *but only at the fixed points !*

Let us now return to our story.

3.1.7 The Closed Strings

No less crucial are the couplings which descend from the hypermultiplets fields arising in the closed string sector of the orbifold. There are twenty such four-component scalar couplings, four controlling the overall shape of the $K3$, while the other sixteen are twisted sector fields, coming one per fixed point of R_{6789} .

A fixed point hypermultiplet's four components split into a triplet ζ of NS-NS fields which transform as a vector of the $SU(2)_R$ group, plus a R-R scalar $\phi^{(0)}$ which is a singlet. As they are closed string fields, they have no group theory charges from the gauge symmetry (D1), or global symmetry (D5, D9) carried by the branes.

They couple into the sigma model as follows. The potential for the 1–1 sector fields b^{AY} arises in an $\mathcal{N}=4$ 1+1 dimensional D–term. In accord with the fact that we have an $SU(2)_R$ symmetry, we can write it in terms of a vector $\boldsymbol{\mu}$.

Using the notation presented in a similar context in ref.[29], the hypermultiplet field b^{AY} is explicitly related to the quaternion:

$$\Psi = \begin{pmatrix} \psi_H^1 & -\psi_H^{2\dagger} \\ \psi_H^2 & \psi_H^{1\dagger} \end{pmatrix} = b, \quad (3.23)$$

which acted on by $SU(2)_L$ and $SU(2)_R$ in the defining representation in the obvious way. We can then write the Lie algebra valued “moment map” vector as

$$\boldsymbol{\mu}^a \equiv \text{Tr} \left[\lambda_V^a \cdot \left\{ \Psi_H^{1\dagger} \boldsymbol{\Sigma} \Psi_H^1 + \Psi_H^{2\dagger} \boldsymbol{\Sigma} \Psi_H^2 \right\} \right] \quad (3.24)$$

where

$$\Psi_H^{1\dagger} = \left(\psi_H^1, -\psi_H^{2\dagger} \right) \quad \text{and} \quad \Psi_H^{2\dagger} = \left(\psi_H^2, \psi_H^{1\dagger} \right) \quad (3.25)$$

are the natural $SU(2)_R$ doublets appearing in the quaternionic form (3.23). The three components, σ^i , of $\boldsymbol{\Sigma}$ are the Pauli matrices acting on the $SU(2)_R$ doublet space. The λ_V^a are generators of the gauge group under which the hypermultiplets are charged.

This vector enters into the Lagrangian coupling to an auxiliary field D . The closed string fields $\boldsymbol{\zeta}$ also couple to the D–auxiliary field, a term arising for each available $U(1)$ factor of the gauge group. In this case, there is one such factor. After integrating D out, we learn that $\boldsymbol{\mu}$ and $\boldsymbol{\zeta}$ couple in Fayet–Iliopoulos fashion

$$\int d^2\sigma \left(\boldsymbol{\mu}^i - \boldsymbol{\zeta} \right)^2, \quad (3.26)$$

where $\boldsymbol{\mu}^i$ means the restriction of $\boldsymbol{\mu}$ in (3.24) to the $U(1)$ subgroup. The non–Abelian parts of $\boldsymbol{\mu}$ simply give D–terms with no FI term contributing^{20,21}. These D–terms are also an important part of the hyperKähler quotient, of course and should not be ignored.

Away from the fixed point, the gauge group is $SO(N)$, and there is no such coupling allowed. This is perfectly acceptable because the closed string twisted sector fields are of course not present in this case.

This coupling is intimately related to the mechanism which is responsible for breaking the $U(16)$ ’s to $SU(16)$ ’s (see ref.[6,34]). In the 5+1 dimensional theory, while the triplet $\boldsymbol{\zeta}$ is involved in this FI term, the final member of the supermultiplet, $\phi^{(0)}$, has Chern–Simons couplings to a six–form “anomaly polynomial” $X^{(6)}$, producing a counterterm

²⁰ That such couplings appear in the physics of D–branes was demonstrated explicitly in calculations in ref.[34].

²¹ See refs.[36,29] for more details of how the coupling works in this context of D1–brane probes.

in the Lagrangian which allows the anomalous $U(1)$ to be canceled in a generalized[37] Green–Schwarz fashion. To complete the job, anomalous gauge transformations are given to $\phi^{(0)}$ which induce a mass term for the $U(1)$, breaking $U(16)$ to $SU(16)$, in a manner generalizing the Dine–Seiberg–Witten[38] mechanism.

3.2. Probing with D1–Branes: The Geometry

3.2.1 Instantons, ALE Spaces, D–Flatness, HyperKähler Quotients and ADHM Data.

At the end of the day, we discover the background fields that the D1–brane probe sees by simply finding the moduli space of possible values which the fields can take, given that we must preserve $(0, 4)$ supersymmetry. This moduli space of vacua, given by $V=0$, where V is the potential, should of course only include (world–sheet) gauge inequivalent field configurations, if we are not to over specify the problem.

A shorthand phrase for this procedure is to perform the “hyperKähler quotient[39]”. Indeed, as first pointed out in refs.[32,34], the physics of D–branes as probes (with this amount of supersymmetry) is isomorphic to certain classic mathematical work by Atiyah, Drinfeld, Hitchin and Manin (ADHM)[40], by Kronheimer[41], and by Kronheimer and Nakajima[42].

This work is extremely relevant to us:

(1) The ADHM construction of instantons essentially constructs the instanton connections as

$$A_\mu = U^\dagger \partial_\mu U, \quad (3.27)$$

where U is defined by

$$\begin{aligned} \mathcal{D}^\dagger U &= 0 \\ U^\dagger U &= \mathbb{I}_2. \end{aligned} \quad (3.28)$$

The operator $\mathcal{D}$ is constructed²² out of what might be called the “ADHM data”, and when it obeys

$$\mathcal{D}^\dagger \mathcal{D} = \Delta \otimes \mathbb{I}_2, \quad (3.29)$$

where Δ is an invertible Hermitian constant matrix, the field strength $F_{\mu\nu}$ of A_μ in (3.27), is self–dual. (U and $\mathcal{D}$ have quaternion values, in the 2×2 defining representation of $SU(2)_R$, and $\mathbb{I}_2$ is the corresponding identity matrix.)

These conditions translate directly into a series of equations for the ADHM data. These equations are precisely the equations (3.18) ensuring $(0, 4)$ supersymmetry of our particular sigma model, (3.17). As first pointed out in ref.[31], $V=0$ for the sigma model (and removing gauge redundancy) gives a space of solutions which is isomorphic to those of ADHM.

²² A most enlightening paper where much of this is made explicit is ref.[43].

In particular, restricting to the massless modes, one finds that the kinetic terms for the current algebra fermions are shifted to[31]:

$$\frac{i}{2} \int d^2\sigma \sum_{i,j} \left\{ \lambda_{+i}^M (\delta_{ij} \partial_- + \partial_- b^\mu A_{\mu ij}) \lambda_{+j}^N \right\}, \quad (3.30)$$

where we have used the x^6, x^7, x^8, x^9 spacetime index μ on our 1–1 field b^{AY} instead of the indices (A, Y) , for clarity.

That the ADMH conditions are the conditions (3.18) on 5–5 and 9–5 fields (couplings here) was shown explicitly in ref.[32].

(ii) Kronheimer’s construction of ALE instantons is also relevant to us. In ref.[41], it was shown that one can construct the ALE spaces using a hyperKähler quotient. The technique shows how to construct the full family of spaces as functions of a set of deformation parameters $\{\zeta\}$. When $\{\zeta=0\}$, one has the singular spaces $\mathbb{R}^4/\Gamma$, where Γ is any discrete subgroup of $SU(2)$ while non-zero ζ gives their smooth deformations into ALE gravitational instantons. “ALE” means that they are flat at infinity, but not Euclidean, due to identifications by Γ at infinity. As Γ has an A–D–E classification[44], so does the family of ALE spaces[45].

In the case in hand, we have $\Gamma=\{1, R_{6789}\}=\mathbb{Z}_2$, and the resolved space is called the Eguchi–Hanson[46] space, controlled by a single deformation parameter a :

$$ds^2 = \left(1 - \left(\frac{a}{r}\right)^4\right)^{-1} dr^2 + \frac{r^4}{4}(\sigma_x^2 + \sigma_y^2) + \frac{r^2}{4} \left(1 - \left(\frac{a}{r}\right)^4\right) \sigma_z^2. \quad (3.31)$$

Here, σ_i are the familiar $SU(2)_R$ –invariant one–forms of $SU(2)$. The metric (3.31) is shown in terms of these to make its $SU(2)_R$ invariance manifest. In Euler coordinates (ϕ, θ, ψ) , we have:

$$\begin{aligned} \sigma_z &= d\psi + \cos\theta d\phi \\ \sigma_y &= -\cos\psi d\theta - \sin\psi \sin\theta d\phi \\ \sigma_x &= \sin\psi d\theta - \cos\psi \sin\theta d\phi \end{aligned} \quad (3.32)$$

giving $\sigma_x^2 + \sigma_y^2 = d\theta^2 + \sin^2\theta d\phi^2$.

It turns out that the physics of D–branes probing orbifold singularities $\mathbb{R}^4/\Gamma$ is isomorphic to Kronheimer’s construction, as shown explicitly for the A–series in refs.[34,36] and for the D and E series in ref.[29]²³. The data specifying the quotient is phrased in terms of a “moment map” μ and some parameters ζ (called the “level set”). In performing the hyperKähler quotient, the equation

$$\mu^i - \zeta = 0 \quad (3.33)$$

²³ See also the appendix of ref.[43] for the explicit example of the derivation of the metric (3.31) with this method.

must be satisfied. This is what we do (among other things, like setting the other D-terms to zero) in setting $V=0$, when we recognize that one of the terms in V is given by the equation (3.26). As mentioned in section 3.1.7, the moment map $\boldsymbol{\mu}$ is simply a combination of 1–1 hypermultiplet fields b^{AY} , while the deformation parameters $\boldsymbol{\zeta}$ are the NS–NS blowup modes coming from the closed string twisted sectors.

(iii) Kronheimer and Nakajima constructed gauge instantons on the ALE spaces by combining the techniques mentioned above in (i) and (ii). The self dual connections A on the ALE space are constructed as

$$A_\mu = U^\dagger \nabla_\mu U, \quad (3.34)$$

where U satisfies similar equations as before in terms of an object $\mathcal{D}$ which this time contains both ADHM type data and moment map type data²⁴. The conditions on $\mathcal{D}$ for self-duality are turn out to be of the same form as equations (3.18), but deformed by $\boldsymbol{\mu}$. The rest of the hyperKähler quotient ($V=0$, etc) includes imposing the blowup condition (3.33) to make the ALE base space.

In our case, the equations which we have written immediately before and after (and including) (3.17) are for the situation where we are away from a fixed point. At a fixed point, we must use the additional multiplets and coupling for the 1–9 sector as discussed in section 3.1.6. The modified conditions for (0,4) supersymmetry are, in the basis we have chosen, of the same form as (3.18), but with extra terms added to include $C_{AA'}^{YM}$. The deformation parameter $\boldsymbol{\zeta}=\boldsymbol{\mu}^i$ is implicit, as $\boldsymbol{\mu}$ is part of the potential for b^{AY} , given in terms of the C ’s.

3.2.2 The Moduli Space of GP Models and Probe Models

So far, we have constructed everything about the D–probe model at the highly symmetric point in the moduli space of GP models where we have gauge symmetry $U(16)_9 \times U(16)_5$ gauge symmetry. This comes from having no Wilson lines for the D9–branes and all of the 32 D5–branes on one fixed point. This gauge symmetry acts as a global symmetry of the 1+1 dimensional probe model, as the all of the fields (except the 1–1 fields) are charged under it, and all of the important couplings (except those descending from the closed string sector) as well.

There is a whole family of models in the GP class, and hence by extension, a whole family of D–probe models. In general, there can be an even number of D5–branes at any fixed point. For m_I such pairs at the I th fixed point, there is enhanced gauge symmetry $U(m_I)_5$ contributed to the gauge group. There are two 5–5 hypermultiplets X_{mn}^{AY} in the

²⁴ The “covariant” derivative ∇ used in (3.34) is defined with a certain connection A^T which is an Abelian connection on the “tautological bundle” on the ALE space. Simply put, the “tautological bundle” is a certain bundle on an ALE space which comes essentially for free. See point (iv) below, for $\mathcal{A}^T$ ’s crucial role in the context of this paper.

$\frac{1}{2}\mathbf{m}_I(\mathbf{m}_I - 1)$ dimensional antisymmetric representation, and a 9–5 hypermultiplet h_{Mn}^A in the $(\mathbf{16}, \mathbf{m}_I)$.

For n'_J D5–branes at a non fixed point J , the gauge group is $USp(2n'_J)$, where n'_J must be a multiple of four, the basic dynamical unit away from a fixed point. There is a 5–5 hyper X_{mn}^{AY} in the $\frac{1}{2}\mathbf{n}'_J(\mathbf{n}'_J - 1)$ antisymmetric representation and a 9–5 hyper h_{Mn}^A in the $(\mathbf{16}, \mathbf{n}'_J)$. This is as it should be for type IB D5–branes in open space, where they should behave like small $Spin(32)/\mathbb{Z}_2$ instantons[35].

The pattern just described is reflected in the structure of the patterns of breaking which can occur with D9–brane Wilson lines, and therefore the possible spectrum of gauge groups and hypermultiplets in the 9–9 sector is identical.

This structure is inherited by the 1+1 dimensional probe model, the 1–9 fields λ_+^M , $\rho^{A'M}$, and $\zeta_\pm^{AM}$, and 1–5 fields $\phi^{A'm}$ and $\chi_\pm^{Am}$, all transform in the fundamentals of the relevant gauge groups. The 9–9, 9–5 and 5–5 couplings are the hypermultiplets with charges given above.

So we see that the various branches of the moduli space includes the physics which we expected to see:

(a) D5–branes are instantons in the D9–brane gauge fields. Away from the fixed points (where 4 D5–branes make a single instanton), the physics is simply that of $Spin(32)/\mathbb{Z}_2$ instantons, as pointed out in ref.[35]. The expectation values of the 9–5 fields control the overall size of the instanton and when their expectation values are zero we have enhanced gauge symmetry $USp(2) \equiv SU(2)$ for each isolated single instanton and $USp(2k)$ when k of them are coincident. Here k is 8, at most because the 32 D5–branes are forced to move in groups of four, as can be deduced from the allowed pattern of Higgs–ing in the 5+1 dimensional model, which translates into an R–symmetry restriction in the sigma model here.

(b) On the fixed points, D5–branes are also instantons, but are now instantons on the resolved space. The small instanton limit there gives enhanced gauge symmetry $U(k)$ for k coincident *half*–instantons (two D5–branes) on the fixed point.

Note however that:

(c) In the absence of D5–branes in the vicinity, and we are just probing the fixed point, our ADHM instanton data is not nearly as complicated as that which specifies the instantons which D5–branes construct. This is because of the absence of an analogue of the 9–5 hypermultiplet fields h_{Mm}^A . The vacuum expectation values of the h control the *size* of the non–Abelian D5–brane instantons. At the fixed point, when we the 1–9 fields (ρ, ζ) and their couplings in the sigma model, there is no analogue of the h fields to give a size modulus to whatever instantons the D1–branes will see coming purely from the D9–branes and the fixed point. So those instantons will be of a special type, geometrically, having no size deformation.

(d) These special instantons will have no size deformations. More correctly, they will have no size parameter independently of the geometry of the resolved fixed point. In other words, their size is set by the same parameter which sets the size of the blowup (see next section). Intuitively, this singles out an (essentially) unique family of instantons associated to the class of spaces which the fixed points resolve to. These “ALE” spaces all possess, due to purely geometrical considerations, a $U(1)$ monopole connection A^T which is naturally associated to the space²⁵. A^T is essentially a monopole field, which is to say that it is an Abelian instanton. In the case (most relevant here) of the Eguchi–Hanson space (3.31), to which our fixed points resolve, the $U(1)$ instanton connection is

$$A^T = -\frac{a^2}{r^2}(d\psi + \cos\theta d\phi). \quad (3.35)$$

We see that the single parameter a controls the size of the Eguchi–Hanson space (3.31) and the (trivial) scale of the Abelian instanton (3.35). This is awfully similar to what we expect from the above comments (c) about the instanton which the D1–branes should see living on the fixed point, and in the absence of D5–branes.

Intuitively, this is clearly the instanton we want²⁶. What we have done by discovering the required form of the sigma model at the fixed point, is to take the Abelian instanton — which is guaranteed to be present for geometrical reasons — and embed it into the D9–brane gauge group. That final step of embedding this instanton, which is otherwise just

²⁵ They are connections on the “tautological bundle” over the ALE space X . The first Chern class of their field strengths F decomposes under Γ into a basis for the $H^2(X, \mathbb{R})$ cohomology of ALE spaces, and the resulting intersection matrix turns out to be isomorphic to the adjacency matrix of the familiar A–D–E Dynkin diagrams. This in turn defines a family of intersecting two–spheres which is the minimal resolution of the orbifold singularity. But this is one beautiful story we must regretfully neglect to tell in full.

²⁶ Mathematically, a proof must work as follows: In order to show that in the absence of 5–9 type couplings, the resulting instantons are simply the canonical Abelian ones, it must be that the matrix U in eqn.(3.34) turns out to be a constant. Then the solution $A=A^T$ results, given the form of the covariant derivative $\nabla=\partial+A^T$. After consulting [42], one sees that the 9–9 hypermultiplets are valued in the endomorphisms $End(W)$ of a certain vector space W , while the 5–9 hypermultiplets are in the homomorphisms $Hom(V, W)$ from W to another vector space V . Roughly, the operator $\mathcal{D}^\dagger$ turns out to be a special map between spaces further defined by taking certain combinations of W, V and $\mathcal{T}$, the tautological bundle. If U is a constant it must mean (given (3.28)) that the kernel of the operator $\mathcal{D}^\dagger$, acting as a map, is trivial. When there is no involvement of the 5–9 fields (and hence no involvement of structures in $Hom(V, W)$), it must be that the structure of the linear maps that $\mathcal{D}^\dagger$ represents simplifies considerably, resulting in the required simple result for U . The resulting moduli of the resulting instantons must have dimension zero. It would be interesting to complete that proof.

surplus to requirements, is not guaranteed at the outset: we have to actually perform it in constructing the consistent theory²⁷. We did so by allowing the 9–9 fields to communicate properly to the 1–1 sector via the new 1–9 fields we discovered.

Hence, we have made contact with section 4.1 of ref.[6]. In the orbifold limit, as we circumnavigate the fixed points we know that the 1–9 fields undergo a monodromy represented by a 32×32 matrix M given by

$$M = \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix} \quad (3.36)$$

where I is the 16×16 identity matrix. This is inherited directly from the choices made in ref.[9]. Normalizing the Abelian instanton A^T such that its monodromy about infinity matches M , gives it $Spin(32)/\mathbb{Z}_2$ instanton number 1, and ensures that it is an instanton without “vector structure”, which is to say that it does not obey Dirac quantization in vector representations of $SO(32)$.

The surviving subgroup of $Spin(32)/\mathbb{Z}_2$ after this embedding is the $U(16)$ which we know that we have from the outset. We have therefore seen from the worldsheet point of view just how the D1–branes see the embedding of the Abelian instanton into the parent gauge group.

3.2.3 The Special Point

As mentioned and studied considerably in section 2, there is a certain special point in the moduli space of GP models which is of great interest. By extension, this point will also define a special 1+1 dimensional sigma model in the class we have built here.

This special point is the placement of two D5–branes on each of the sixteen fixed points in the orbifold $K3$. The model is connected to the situation where there is no gauge group. Considering the six dimensional anomaly equation:

$$n_H - n_V = 244 + 29n_T, \quad (3.37)$$

where n_H, n_V and n_T represent the number of hyper–, vector– and tensor– multiplets, respectively, this means that there are 244 hypermultiplets from the open and closed string sectors, as there are no extra tensor multiplets in the spectrum beyond the standard one on the supergravity multiplet²⁸.

Let us see how the various hypermultiplet contributions arise at this special point. Placing a single D5–brane pair at each of the sixteen fixed points (and turning on appropriate Wilson lines on the torus) we obtain gauge group $U(1)_9^{16} \times U(1)_9^{16}$. At this stage $n_V=32$

²⁷ Contrast this with the case of the type II theories (*e.g.* ref.[29] in the same situation where the Abelian connections on the ALE spaces are certainly there, but play no role in a stringy gauge bundle.

²⁸ See, however refs.[16,17] for orientifold models similar to the GP models which have extra tensors.

and therefore $n_H=276$. By examining the details of the GP spectrum given in section 3.2.2, we see that there are no 5–5 or 9–9 fields remaining, but there are 16 5–9 fields coming from each of the sixteen fixed points making 256 9–5 fields. The remaining 20 are the $K3$ deformation hypermultiplets from the closed string sector, 16 of them coming from the fixed points as twisted sector fields. As pointed out in ref.[6], 16 of the $U(1)$ ’s give rise to anomalies which are canceled at one loop by a generalization of the Green–Schwarz mechanism[10] which simultaneously lifts the gauge symmetries by a generalization of the Dine–Seiberg–Witten mechanism[38]. 16 R–R components of the closed string twist fields are involved in this process, their 16 NS–NS triplet partners, ζ_i , remaining to control the blowup of the $K3$, as discussed in sections 3.1.7 and 3.2.1. Giving vacuum expectation values to these will therefore remove the 16 anomalous $U(1)$ ’s. The remaining 16 $U(1)$ ’s can be Higgsed away, using up a 9–5 field at each fixed point²⁹.

In the end, therefore, we see that the 244 hypermultiplets of the spectrum of the special GP point are made up of 4 closed string hypers, controlling the global geometry of the $K3$, and 15 9–5 fields coming from each fixed point. (The size of the vacuum expectation values of the 16 closed string twist fields are still adjustable in order to control the $K3$ shape, of course.) These open string hypermultiplets are also pure geometry: They are coordinates on the Higgs branch (of the six dimensional field theory) of the $SO(32)$ (half–) instanton moduli space, representing the deformations away from the zero size limit. As there are no 5–5 fields remaining, there is no branch representing the motion of the point–like half–instantons; they are stuck on the fixed points and all they can do is grow and shrink. There are no moduli for the Abelian instanton of the 9–9 sector living at the fixed point, as we have seen explicitly from equations (3.31) and (3.35), and the discussion in section 3.2.2.

This is also confirmed by the index theorem computations in ref.[6], where it was computed that on the manifold $K3$, the dimension of the moduli space $\mathcal{M}_k$ of k $Spin(32)/\mathbb{Z}_2$ instantons which break the gauge group completely, with corrections included to take into account the possibility of fractional instantons, is

$$\dim \mathcal{M}_k = 120(k - 1). \quad (3.38)$$

From this, ref.[6] saw that for $k=1$, the case of the instanton inherent to the fixed point, we get the answer zero, confirming that there are no moduli for that instanton.

Here, we see that this formula confirms everything else we have said above, since with the half–instanton included, we have 3/2 of an instanton at each point. Putting this into the formula, we get that there is a 60 dimensional moduli space associated with this point. Dividing by four to get the amount of hypermultiplets, this corresponds to the 15 hypermultiplets which we found above.

²⁹ In the T_{6789} –model, the 5–5 $U(1)$ ’s are Higgs’d and the 9–9 $U(1)$ ’s are Dine–Seiberg–Witten’d.

In this way, we understand — from many complementary points of view — just what the one dimensional probe of N coincident D1–branes sees as it moves around the interior of the $K3$ of our compactification. In particular, there is $3/2$ of an instanton stuck at each fixed point of the orbifold $K3$.

3.3. Probing with D0–branes and D5–branes.

3.3.1 D0–brane probes

Now we are in a position to consider the N D0–brane probe 0+1 dimensional theory. There is not much to do, in terms of constructing a Lagrangian, as it is obtained by simply dimensionally reducing the one we have constructed in the previous subsection.

Thinking of this dimensional reduction as being performed on the x^5 circle, we must take into account the possibility of introducing Wilson lines for the gauge field A_μ in the process. This results in an extra massless parameter X^5 in the quantum mechanics representing the position of the D0–branes along the dual x^5 direction, which is the line interval $\mathcal{I}$. Simultaneously, from the point of view of the D9– and D5–branes’ six dimensional gauge fields, there is a reduction to five dimensions with a Wilson line.

For the D9–branes, the particular Wilson line we are interested in is the one which places the resulting D8–branes (in the type IIA picture) symmetrically at the ends of the interval $\mathcal{I}$. Meanwhile, for the D5–branes, we choose Wilson lines which place the 32 D4–branes in the configuration discussed in section 2.3.2. The N D0–brane probe therefore sees 1 D4–brane plus 1 O4–plane at each fixed point of the $K3$ at each end of the interval, equivalent to $3/4$ of an instanton.

(Perhaps less confusingly, we can equivalently think of this as the same $K3$ as in the type IB model, but now every point on it has the content of a line interval $\mathcal{I}$. In this way, we see that there is still the $3/2$ instanton at each fixed point, but it further subdivided into two parts placed at each end of $\mathcal{I}$.)

3.3.2 D5–brane probes

A six dimensional field theory may be obtained from our 1+1 dimensional probe models by T_{6789} –duality. This turns our family of N coincident D1–brane probes into a family of N coincident D5’–brane probes, in the notation of section 2 (see Table 2). They are wrapped around the $K3$ of type IB, and give rise to a dual family of D1–brane probes. By the self–duality of the model, these dual D1–probes have the same physics of the D1–brane probes we discussed in section 3.1 and 3.2. They inherited this physics from the wrapping of the D5’–branes, and hence we indirectly know some of the physics of the compactified N D5’–branes.

We can say a few things directly about the 5+1 dimensional world–volume theory of the wrapped branes which will be useful in a later discussion.

The background D5– and D9–branes produce the same family of global symmetries as they did in the 1+1 dimensional model, supplying charges for the 5′–5 and 5′–9 fields. The 5–5, 9–9 and 5–9 fields enter again as parameters.

We have the usual $\mathcal{N}=1$ supersymmetric Yang–Mills terms in the Lagrangian, in terms of gauge potential A^μ , arising in the 5′–5′ sector from $(x^0, x^5, x^6, x^7, x^8, x^9)$ excitations, and hypermultiplets coming from the 5′–5′ sector with excitations in the (x^1, x^2, x^3, x^4) directions. The gauge group is $U(N)$, the same as for the 1+1 dimensional model at a fixed point. To get the sector where it is $SO(N)$, we simply introduce the Wilson lines which are dual to moving the D1–brane probes away from them as described in subsection 3.2.

Crucially, we also have terms of the form

$$\mu_5 \int d^6y H^{(7)} \cdot H^{(7)} + A^{(2)} \wedge (R \wedge R + F \wedge F), \quad (3.39)$$

where R is the Ricci two–form of $K3$ and F is the field strength of the gauge field A^μ ($\mu=\{0, 5, 6, 7, 8, 9\}$). $H^{(7)}$ is the field strength of the R–R potential $A^{(6)}$ (to which the D5–branes couple “electrically” with charge μ_5) and $A^{(2)}$ is its ten dimensional dual.

Satisfying the resulting field equations requires that due to the presence of the 16 sources of curvature ($K3$ ’s fixed points), we have to have 16 gauge instanton sources. These are of course, string–like objects in the world volume theory, which are known to be D1–branes[33], objects which are sources of $A^{(2)}$.

We can say precisely what the nature of these 16 D1–branes living in the D5–branes must be. First of all, they are identical in structure, as the 16 fixed points are all of the same type. They are necessarily transverse to the $K3$ (and hence the coordinates where the gauge instanton they carry is located), which puts them in the (x^0, x^5) directions. Furthermore, they must have precisely the same global symmetries on their world–sheet action as the D5–branes’ world–volume possesses. *This identifies them with exactly the family of D1–branes we studied in sections 3.1 and 3.2.*

The crucial lesson of this discussion, which we will revisit later on, is as follows: There is a moduli space of $K3$ compactifications of the 5+1 dimensional theory associated with the N D5′–branes, which encodes the data of the T_{6789} –dual $K3$ compactification of the type IB theory. We start with N D5′–branes in flat space, which has a $U(N)$ gauge symmetry and a $Spin(32)/\mathbb{Z}_2$ global symmetry. We then compactify them on a $K3$. The data associated to compactifying type IB on the dual $K3$ is encoded in the world–volume theory by the patterns of breakings of the global symmetry and the pattern of induced couplings which arise. A family of stringy states are seen to exist in the model. The family of strings which can arise is also parameterized by the patterns of breakings of the global symmetries, and the couplings which can arise in the 1+1 dimensional world–sheet theory. We have characterized such string sigma models in sections 3.1 and 3.2.

4. Some (Partial) Matrix Theory Proposals

We are now in a position to suggest a role for the D-brane theories we have been discussing in the context of Matrix theory. Let us do this step by step:

4.1. Matrix Theory of M-theory on $X \times \mathcal{I}$.

The 1+1 dimensional models which we constructed in section 3 can be used as the starting point for a definition of a matrix theory representation of M-theory compactified on the Eguchi–Hanson space X times a line interval $\mathcal{I}_5$, of length L_M^5 . By equations (2.5) and (2.6), we can deduce precisely what string theory lengths, L_{IA}^5 and L_{IB}^5 , we must use to define the model. In particular, the model should be defined by placing our 1+1 dimensional model —at the special point— on a circle of radius L_{IB}^5 and taking the large N limit. As N is correlated (by T_5 -duality) to D0-brane number, we expect that this model —when accompanied by the large $\lambda_{IA}=R_M^{10}$ (in 11D metric) limit— should give a matrix theory definition of the compactification in the infinite momentum frame. We expect that the problems noticed at finite N for Matrix theories on such surfaces in ref.[47] will disappear at large N , as suggested for example in ref.[48]. This expectation should be accompanied by some hope also, as the physics is rather less constrained by supersymmetry than before[12]: we only have eight supercharges.

4.2. Matrix Theory of the $E_8 \times E_8$ Heterotic String on $X \times \mathcal{I}$.

One interesting limit of the previous compactification is of course the $L_M^5 \rightarrow 0$ limit where we should recover the $E_8 \times E_8$ heterotic string compactified on the Eguchi–Hanson space X , with $3/2$ of an E_8 instanton³⁰.

In the context of our matrix model, this limit is defined by starting with the large (N, R_M^{10}) limit of our 1+1 dimensional model — at the special point — of section 3. It is the decompactified x^5 limit of the model above, referring to the base space where the 1+1 dimensional theory lives. It is natural to infer therefore that the 1+1 dimensional model flows in this limit (which is its IR limit) to a conformal field theory of the heterotic string on X , thus realizing the duality of the M-theory configuration to the heterotic one as a flow³¹. This would be a complicated example of the work of Motl[50], and of Banks and Seiberg[51], and of Dijkgraaf, Verlinde and Verlinde[28]. There a “matrix string theory” of the light cone type IIA string was recovered from Matrix theory as a flow to an S_N orbifold conformal field theory from a 1+1 dimensional model arising essentially from a collection

³⁰ Matrix theories of E_8 instantons have been presented in the literature lately[49], in work independent of the present paper.

³¹ The issue of realizing duality as a flow from D-brane world volume theories was discussed a while ago in the final section of ref.[29].

of N D1-brane probes' common world-volume. In that case, with D1-branes in flat space, the moduli space (and hence the target space of the sigma model) was $(\mathbb{R}^8)^N/S_N$ (where S_N is the group of permutations of N objects), which ultimately defined the conformal field theory. (The uncompactified heterotic matrix theory was studied in refs.[52], and some heterotic toroidal compactifications in refs.[53,54].) In this case, the moduli space of the N D1-brane system needs to be carefully examined (we have not done it fully here) to discover the precise nature of the conformal field theory this new matrix string theory realizes. On general grounds, with everything above as motivation, we expect of course to define the $E_8 \times E_8$ heterotic string on $X \times \mathcal{I}$ in this way.

4.3. Matrix Theory of M-theory on $K3 \times \mathcal{I}$ and its Heterotic Limits.

Next, we come to ask about the definition of M-theory on $K3 \times \mathcal{I}$. Restricting ourselves to the point we know best —the special point— we can anticipate that a six dimensional theory with $(1,0)$ supersymmetry is involved in defining this model.

At least part of this theory is motivated in terms of the world volume theory of N D5'-branes compactified on $\widetilde{K3} \times S_5^1$, along the lines described in section 3.3.2. This probe theory should have some interesting properties, and cannot be the naive super Yang-Mills theory, for the same reasons as discussed in refs.[14,15,55] in the case of the $(0,2)$ situation.

What properties must we require this compactified $(0,1)$ theory to have? One of the motivations of this paper is to point out that at least at a special point in moduli space, *the theory must reproduce the special duality properties of the DMW point*: There should be dual realizations of the $E_8 \times E_8$ heterotic string theory compactified on $K3$ with the $(12,12)$ arrangement of instantons³².

This works as follows: Heterotic/heterotic duality has its M-theory origins in terms of electromagnetic duality exchanging M2- and M5-branes, as first suggested in ref.[4] and

³² A crucial phenomenon we observe here is the fact that the strong coupling limit of going to M-theory mixes up the role of the various hypermultiplets in our spectrum in a way which would not have been consistent with string theory if we had remained at weak coupling. As pointed out last section, of the 244 hypermultiplets not involved in removing various $U(1)$'s, 4 of them are closed string fields, and the rest are 16 copies of 15 (5-9) open string fields, associated to the $3/2$ instantons at each fixed point. By the time we get to M-theory on $K3 \times \mathcal{I}_5$, we have (of course) the same number of hypermultiplets, but their origin is different. 112 hypermultiplets come from placing 12 E_8 instantons on the $K3$ and breaking the E_8 gauge symmetry associated with the end of the interval completely, and another 112 from the other end of the interval. The remaining 20 hypermultiplets come from the family of $K3$ deformations. So we see that at each fixed point, a 5-9 field becomes associated at strong coupling with a deformation of $K3$, while the other 14 are the $224/16$ E_8 instanton deformations.

made explicit in the present paper. In sections 2.6 and 2.7, we showed how to reduce the problem of studying the details of heterotic/heterotic duality in M-theory to one of studying T-duality in the type IA model, and a related T-duality in the type IB model, where in that context its relevance was shown in ref.[6]. So the existence of heterotic/heterotic duality is guaranteed by the existence of a T_{456789} -duality property which exchanges the two type IA models (realizing electric-magnetic duality in M-theory), and hence by a T_{6789} -duality which exchanges the two type IB models.

This translates therefore in this matrix theory context to the statement that (at this special point) heterotic/heterotic duality is ensured by asking that the compactified $(0, 1)$ six dimensional theory has as a basic property a T_{6789} -duality symmetry, giving us the same theory back. (In this limit where we are studying everything explicitly as orbifolds, such a duality on the base space makes the usual good sense. Beyond the orbifold limit, it is not clear what to do.)

This property, as stressed by Seiberg[30], rules out the $(0, 1)$ fixed point field theories as candidates, and rules in the “little” string theories as possible candidates for our matrix definition. This is encouraging, as we saw in section 3.3.2 that we identified excitations in the compactified model which was a particular family of stringy instantons. We also deduced that part of their definition was given by the 1+1 dimensional sigma model of earlier subsections.

The 1+1 dimensional sigma model will play two crucial roles in this situation then. Most naturally, it will flow to a conformal field theory and define the heterotic matrix theory of the $E_8 \times E_8$ heterotic string compactified on $X \times \mathcal{I}$, as described in section 4.2. Alternatively however, there is a limit where the model flows to a point where gravity decouples, defining a matrix theory of a “little $E_8 \times E_8$ heterotic string” in six dimensions, compactified on the base space $\widetilde{K3} \times S_5^1$. Such a dual role fits all of the data and is conceptually satisfying.

In addition to seeing the stringy excitations of the compactified little string model, we also saw (in section 3.3.2) clues about exactly what it means to compactify such a theory on $K3 \times S_5^1$ for the purposes of our definition. We saw that the moduli space of compactifications of the N D5'-brane theory encodes the moduli space of choices to be made about the arrangements of D5-branes and D9-branes in the GP models, which is a particular case of constructing a dual heterotic compactification: This amounts to choosing the gauge bundle and instantons carefully in the presence of the $K3$, breaking the $Spin(32)/\mathbb{Z}_2$ gauge symmetry accordingly, as described in great detail earlier. From the point of view of the little string theory, we are making the same choices about breaking a $Spin(32)/\mathbb{Z}_2$ global symmetry. This seems at odds with the fact that we concluded that we had the $E_8 \times E_8$ little string theory earlier, but recall that precisely at this special point in moduli space, the “big” $Spin(32)/\mathbb{Z}_2$ and $E_8 \times E_8$ heterotic strings are T_5 -dual. That T_5 -duality must be inherited by the compactified little string theories as well, and is another requirement for our compactified $(0, 1)$ theory.

So in short, we see that the moduli space of compactifications of the little heterotic string theory is indeed just that which we want, in order to reproduce the physics of the compactified (big) heterotic string at the special DMW point.

5. Beyond the Special Point

As the theory at the special point is defined in terms of N D5-branes of type IB, we have implicitly made a statement about the $(0,1)$ theory associated to the $Spin(32)/\mathbb{Z}_2$ global symmetry and its compactifications. This is strengthened by the fact that strong/weak coupling duality takes us directly to a family of N $Spin(32)/\mathbb{Z}_2$ heterotic fivebranes, making direct contact with the construction of Seiberg[30]. It so happens that the particular point in moduli space we have studied has an accidental T_5 -duality symmetry to a similarly compactified $E_8 \times E_8$ little string theory. Away from the special point however, one expects that it is the latter which will define the full moduli space of $E_8 \times E_8$ heterotic string $K3$ compactifications³³, although the details of the prescription are harder to motivate. This is because we no longer have the luxury of a complete D-brane construction to help us build sigma models for study, and we also do not have as much control over the properties of $K3$ away from the orbifold limits.

However, it is a highly suggestive that such a definition may not be far from the mark. If correct, we would also gain a definition of the $SO(32)$ heterotic string on $K3$ with the standard embedding, as this is the $(16, 8)$ $E_8 \times E_8$ model[56].

6. Discussion and Conclusion

6.1. Summary

This paper was divided into two main parts. First, we did a thorough examination of some of the properties of the heterotic/heterotic duality of ref.[4]. This was made possible by the fact that it has a realization as a type IB orientifold model[9], as pointed out in ref.[6].

We exploited this fact by turning this orientifold definition into a type IA one, and noticed that precisely at the point of interest, we can take the limit where the model becomes an M-theory compactification, and from there made contact again with the heterotic model. Along the way, we sharpened our understanding of the heterotic/heterotic dual phenomenon and its links to M-theory's electric-magnetic duality properties.

³³ Some alternative comments about the matrix theory for the full moduli space of the heterotic string on $K3$ are made in the conclusions of ref.[54]. I am grateful to S. Govindarajan for pointing this out.

Such an exploration is interesting in its own right and we could have stopped there, but we went further to the second half of the paper. Given that we have a family of orientifold models which connects so nicely the heterotic string on $K3$ to M-theory, on one hand, and to type I theory on the other, it is natural to see whether we might make some headway in defining what the nature of a Matrix theory representation of this M-theory compactification might be. Indeed, if we cannot say anything else about the theory (due to its admittedly low amount of supersymmetry, and its points of spacetime curvature which make D-brane matrix definitions problematic, at least at finite N [47]) we can at least try to constrain its properties by requiring it to reproduce some of the intricate properties of the heterotic/heterotic situation.

This is what we set out to do in the second half of the paper. Along the way, we derived a number of interesting observations which have wider application. The 1+1 dimensional probe model of the D1-branes moving in the orientifold model background was particularly interesting, generalizing the work of refs.[31,34,32] to situations where the gauge bundles of the heterotic compactification are of just the right type to realize the DMW duality.

Now holding some definite models with which to make a Matrix theory proposal, we went on to discuss the properties of the various matrix models which would define the various parts of the theory. We completed the proposal with the conclusion that the central role is played by the “little heterotic string theories”, where we can (at the special point) make a precise statement about the nature of the base space we are to compactify them on, and the identification of the moduli space of those compactifications with those of the usual, “big” heterotic strings. It is natural to suppose that away from the special point, the compactified (0,1) little string theories should be used to define the matrix theory representations of the wider class of heterotic compactifications too, although there is less evidence to support this conjecture.

6.2. New Directions

There is a wealth of interesting issues uncovered here. In particular, more study of the 1+1 dimensional model presented here should be carried out. Part of the moduli space theory contains the physics of E_8 instantons on the Eguchi–Hanson space, which is interesting in its own right. Furthermore, the model can be used to define (by RG flow) an interesting conformal field theory of the matrix heterotic string in such a background, and the details of the moduli space needs to be pinned down in order to identify correctly the properties of the candidate conformal field theory. Ultimately, deformation of the conformal field theory should help construct (by analogy with refs.[50],[28])) the full string field theory perturbation series for heterotic strings in that background.

In another limit this 1+1 model will actually flow to the little heterotic string itself. This is a more careful limit where gravity decouples, and the resulting string is forced to live in the six dimensional situation described above, now compactified on $\widetilde{K3} \times S^1_5$. The

determination of the moduli space of the model will be important for this application too. We have only completed a partial study in this paper.

It is interesting to consider also the case of other orientifold models. Generically, most such models will not satisfy our criteria of section 2.3.2 for being able to be taken properly to M-theory³⁴ There is at least one example of a model which passes the test. The “ $\mathbb{Z}_4^A$ ” model of ref.[16] has an arrangement of D4-branes which allows complete local cancellation of dilaton charge, thereby promising an interesting dual realization. A particularly interesting property of that model is the fact that it has $n_T=4$; there are four extra tensor multiplets in the spectrum. As pointed out in ref.[57], there is strong evidence that it is related to the $(10, 10) + 4$ M-theory $K3_M \times \mathcal{I}$ compactification, having four M5-branes in the interior of $\mathcal{I}$. (It was also argued³⁵ in ref.[57] to be a limit of F-theory on a Calabi–Yau manifold with Hodge numbers $(127, 7)$, which has since been shown to be true[59].) Such a model is therefore not completely to be understood as a heterotic string compactification. Because of its special properties as an orientifold model however, and in the light of this paper, it might be used as a test-bed for a matrix theory representation of the $(10, 10) + 4$ M-theory compactification. Much the same procedures as carried out in this paper would be relevant to such a study. An obvious guess at the result would be that the $E_8 \times E_8$ little string theory should be compactified on an appropriate dual surface (which is meaningful because in that model $K3$ is realized as a $\mathbb{Z}_4$ orbifold of a torus) with some extra features to give the four extra fivebranes. Presumably these features involve tuning four of the 24 E_8 constituent instantons such that they move out onto the branch of their moduli space where a tensor arises in exchange for 29 hypermultiplets.

There are many avenues of interest to pursue using the constructions of this paper as a starting point. Hopefully such further research will help to further clarify the properties of Matrix theory and the little string theories.

³⁴ I am grateful to R. C. Myers for discussions on the interesting issue of whether there are other ways of controlling the limit of six dimensional type IA orientifold models in such a way as to get M-theory configurations.


³⁵ In that paper, similar statements also were made for the $\mathbb{Z}_6^A$ orientifold model of ref.[16]. This has since been shown by E. Gimon[58] to be incorrect. It is not possible to arrange the D5-branes in the way suggested there in order to cancel the dilaton charges locally, and therefore more care is needed with this model.

Acknowledgments

CVJ was supported in part by family, friends and music. Thanks to the Aspen Center for Physics, where most of this work was carried out, and the organizers of the Duality workshop (July 13th to August 3rd 1997), for an inspiring meeting. Thanks also to the organizers of the International Symposium on ‘New Trends in Subatomic Physics’, (National Taiwan University, August 5th to 13th 1997), for an interesting and timely gathering, and where most of the rest of this work was carried out in the remarkably fine surroundings of the Grand Hotel, Taipei.

CVJ would like to thank Ofer Aharony, Eric Gimon, Ami Hanany, Robert Myers, Hiroshi Ooguri, Eva Silverstein and Edward Witten for conversations.

Travel support was obtained under Department of Energy grant # DE-FC02-91ER75661.



References

- [1] E. Witten, ‘Five-branes And M-Theory On An Orbifold’, Nucl. Phys. **B463** (1996) 383, hep-th/9512219.
- [2] R. Güven, ‘Black p -Brane Solutions of $D=11$ Supergravity theory’, Phys. Lett. **B276** (1992) 49.
- [3] E. Bergshoeff, E. Sezgin and P. K. Townsend, ‘Supermembranes and Eleven Dimensional Supergravity’, Phys. Lett. **B189** (1987) 75;
M. J. Duff and K. S. Stelle, ‘Multimembrane solutions of $D=11$ Supergravity’, Phys. Lett. **B253** (1991) 113.
- [4] Duff, Minasian and Witten, ‘Evidence for Heterotic/Heterotic Duality’, Nucl. Phys. **B465** (1996) 413, hep-th/9601036.
- [5] M. J. Duff, ‘Strong/Weak Coupling Duality from the Dual String’, Nucl. Phys. **B442** (1995) 47, hep-th/9501030.;
M. J. Duff, ‘Putting string/string duality to the test’, Nucl. Phys. **B436** (1995) 507, hep-th/9406198.
- [6] M. Berkooz, R. G. Leigh, J. Polchinski, J. Schwarz, N. Seiberg and E. Witten, ‘Anomalies, Dualities, and Topology of $D=6$ $N=1$ Superstring Vacua’, Nucl. Phys. **B475** (1996) 115, hep-th/9605184.
- [7] J. Dai, R. G. Leigh and J. Polchinski, ‘New Connections Between String Theories’, Mod. Phys. Lett. **A4** (1989) 2073;
P. Hořava, ‘Background Duality of Open String Models’, Phys. Lett. **B231** (1989) 251;
R. G. Leigh, ‘Dirac–Born–Infeld Action from Dirichlet Sigma Model’, Mod. Phys. Lett. **A4** (1989) 2767;
J. Polchinski, ‘Combinatorics Of Boundaries in String Theory’, Phys. Rev. **D50** (1994) 6041, hep-th/9407031.
- [8] A. Sagnotti, in ‘Non–Perturbative Quantum Field Theory’, Eds. G. Mack *et. al.* (Pergammon Press, 1988), p.521;
V. Periwal, unpublished;
J. Govaerts, Phys. Lett. **B220** (1989) 77;
P. Horáva, ‘Strings on World Sheet Orbifolds’, Nucl. Phys. **B327** (1989) 461.
- [9] E. G. Gimon and J. Polchinski, ‘Consistency Conditions of Orientifolds and D -Manifolds’, Phys. Rev. **D54** (1996) 1667, hep-th/9601038.
- [10] M. Bianchi and A. Sagnotti, ‘Twist Symmetry and Open String Wilson Lines’ Nucl. Phys. **B361** (1991) 519.

- [11] G. Aldazabal, A. Font, L.E. Ibanez, F. Quevedo, ‘*Heterotic/Heterotic Duality in $D=6,4$* ’, Phys.Lett. **B380** (1996) 33, hep-th/9602097.
- [12] T. Banks, W. Fischler, S. Shenker and L. Susskind, ‘*M–Theory As A Matrix Model: A Conjecture*’, Phys. Rev. **D55** (1997) 5112, hep-th/9610043.
- [13] W. Taylor, ‘*D–Brane field theory on compact spaces*’, Phys. Lett. **B394** (1997) 283, hep-th/9611042;
O. J. Ganor, S. Ramgoolam, W. Taylor, ‘*Branes, Fluxes and Duality in M(atrix)-Theory*’, Nucl. Phys. **B492** (1997) 191, hep-th/9611202.
- [14] M. Rozali, ‘*Matrix Theory and U–Duality in Seven Dimensions*’, Phys. Lett. **B400** (1997) 260, hep-th/9702136.
- [15] M. Berkooz, N. Seiberg and M. Rozali, ‘*Matrix Description of M-theory on T^4 and T^5* ’, Phys. Lett. **B408** (1997) 105, hep-th/9704089.
- [16] E. G. Gimon and C. V. Johnson, ‘*K3 Orientifolds*’, Nucl. Phys. **B477** (1996) 715, hep-th/9604129.
- [17] A. Dahbolkar and J. Park, ‘*Strings on Orientifolds*’, Nucl. Phys. **B477** (1996) 701, hep-th/9604178.
- [18] P. Horava and E. Witten, ‘*Heterotic and Type I String Dynamics from Eleven Dimensions*’, Nucl. Phys. **B460** (1996) 506, hep-th/9510209.
- [19] J. Polchinski, ‘*TASI Lectures on D-Branes*’, hep-th/9611050.
- [20] J. Polchinski, S. Chaudhuri and C. V. Johnson, ‘*Notes on D–Branes*’, hep-th/9602052.
- [21] E. Bergshoeff, M. de Roo, M. B. Green, G. Papadopoulos and P. K. Townsend, ‘*Duality of Type II 7-branes and 8-branes*’, Nucl.Phys. **B470** (1996) 113, hep-th/9601150.
- [22] L. Romans, ‘*Massive $N=2a$ Supergravity In Ten-Dimensions*’, Phys. Lett. **B169** (1986) 374.
- [23] J. Polchinski, ‘*Dirichlet Branes and Ramond–Ramond Charges in String Theory*’, Phys. Rev. Lett. **75** (1995) hep-th/9510017.
- [24] E. Witten, ‘*String Theory Dynamics in Various Dimensions*’, Nucl. Phys. **B443** (1995) 85, hep-th/9503124.
- [25] J. Polchinski and E. Witten, ‘*Evidence for Heterotic - Type I String Duality*’, Nucl. Phys. **B460** (1996) 525, hep-th/9510169.
- [26] C. M. Hull, ‘*String–String Duality in Ten Dimensions*’, Phys. Lett. **B357** (1995) 545, hep-th/9506194.
- [27] S.-J. Rey, ‘*Axionic String Instantons and Their Low–Energy Implications*’, in ‘*Superstrings and Particle Theory*’, proceedings, eds. L. Clavelli and B. Harms, World

- Scientific, 1990;
- S.-J. Rey, ‘*The Confining Phase of Superstrings and Axionic Strings*’, Phys. Rev. **D43** **1991** 526;
- A. Strominger, ‘*Heterotic Solitons*’ Nucl. Phys. **B343**, (1990) 167; *Erratum: ibid.*, **353** (1991) 565.
- [28] R. Dijkgraaf, E. Verlinde, H. Verlinde, ‘*Matrix String Theory*’, Nucl. Phys. **B500** (1997) 43, hep-th/9703030.
- [29] C. V. Johnson and R. C. Myers, ‘*Aspects of Type IIB Theory on Asymptotically Locally Euclidean Spaces*’, Phys. Rev. **D55** (1997) 6382, hep-th/9610140.
- [30] N. Seiberg, ‘*Matrix Description of M-theory on T^5 and T^5/Z_2* ’, Phys. Lett. **B408** (1997) 98, hep-th/9705221.
- [31] E. Witten, ‘*Sigma Models and the ADHM Construction of Instantons*’, J. Geom. Phys. **15** (1995) 215, hep-th/9410052.
- [32] M. R. Douglas, ‘*Gauge Fields and D-Branes*’, hep-th/9604198.
- [33] M. R. Douglas, ‘*Branes within Branes*’, hep-th/9512077.
- [34] M. R. Douglas and G. Moore, ‘*D-Branes, Quivers and ALE Instantons*’, hep-th/9603167.
- [35] E. Witten, ‘*Small Instantons in String Theory*’, Nucl. Phys. **B460** (1996) 541, hep-th/9511030.
- [36] J. Polchinski, ‘*Tensors From K3 Orientifolds*’, hep-th/9606165.
- [37] A. Sagnotti, ‘*A Note on the Green - Schwarz Mechanism in Open - String Theories*’, Phys. Lett. **B294** (1992) 196, hep-th/9210127.
- [38] M. Dine, N. Seiberg and E. Witten, ‘*Fayet-Iliopoulos Terms in String Theory*’, Nucl. Phys. **B289** (1987) 589.
- [39] N. J. Hitchin, A. Karlhede, U. Lindström and M. Roček, ‘*Hyper-Kähler Metrics and Supersymmetry*’, Comm. Math. Phys. **108** (1987) 535.
- [40] M. F. Atiyah, V. Drinfeld, N. J. Hitchin and Y. I. Manin, ‘*Construction of Instantons*’ Phys. Lett. **A65** (1978) 185.
- [41] P. B. Kronheimer, ‘*The Construction of ALE Spaces as Hyper-Kähler Quotients*’, J. Diff. Geom. **29** (1989) 665.
- [42] P. B. Kronheimer and H. Nakajima, ‘*Yang-Mills Instantons on ALE Gravitational Instantons*’, Math. Ann. **288** (1990) 263.
- [43] M. Bianchi, F. Fucito, G. Rossi, and M. Martinelli, ‘*Explicit Construction of Yang-Mills Instantons on ALE Spaces*’, Nucl. Phys. **B473** (1996) 367, hep-th/9601162.

- [44] J. McKay, ‘*Graphs, Singularities and Finite Groups*’, Proc. Symp. Pure. Math. **37** (1980) 183, Providence, RI; Amer. Math. Soc.
- [45] N. J. Hitchin, ‘*Polygons and Gravitons*’, Math. Proc. Camb. Phil. Soc. **85** (1979) 465.
- [46] T. Eguchi and A. J. Hanson, ‘*Asymptotically Flat Self-Dual Solutions to Euclidean Gravity*’, Phys. Lett. **B74** (1978) 249.
- [47] M. R. Douglas, H. Ooguri, S. H. Shenker, ‘*Issues in M(atric) Theory Compactification*’, Phys.Lett. **B402** (1997) 36, hep-th/9702203;
M. R. Douglas, H. Ooguri ‘*Why Matrix Theory is Hard*’, hep-th/9710178.
- [48] W. Fischler, A. Rajaraman, ‘*M(atric) String Theory on K3*’, hep-th/9704123.
- [49] D. Lowe, ‘ *$E_8 \times E_8$ Instantons in Matrix Theory*’, hep-th/9709015;
O. Aharony, M. Berkooz, S. Kachru, and E. Silverstein, ‘*Matrix Description of (1,0) Theories in Six Dimensions*’, hep-th/9709118.
- [50] L. Motl, ‘*Proposals on Non-Perturbative Superstring Interactions*’, hep-th/9701025.
- [51] T. Banks and N. Seiberg, ‘*Strings from Matrices*’, hep-th/9702187.
- [52] U. H. Danielsson, G. Ferretti, ‘*The Heterotic Life of the D-particle*’, Int. J. Mod. Phys. **A12** (1997) 4581, hep-th/9610082;
S. Kachru and E. Silverstein, ‘*On Gauge Bosons in the Matrix Model Approach to M Theory*’, Phys. Lett. **B396** (1997) 70, hep-th/9612162;
L. Motl, ‘*Quaternions and M(atric) theory in spaces with boundaries*’, hep-th/9612198;

N. Kim and S.-J. Rey, ‘*M(atric) Theory on an Orbifold and Twisted Membrane*’, Nucl. Phys. **B504** (1997) 189;
D. Lowe, ‘*Bound States of Type I’ D-particles and Enhanced Gauge Symmetry*’, Nucl. Phys. **B501** (1997) 134, hep-th/9702006;
T. Banks, N. Seiberg, E. Silverstein, ‘*Zero and One-dimensional Probes with N=8 Supersymmetry*’, Phys.Lett. **B401** (1997) 30, hep-th/9703052;
T. Banks and L. Motl, ‘*Heterotic Strings from Matrices*’, hep-th/9703218;
D. Lowe, ‘*Heterotic Matrix String Theory*’, Phys. Lett. **B403** (1997) 243, hep-th/9704041;
S.-J. Rey, ‘*Heterotic M(atric) Strings and Their Interactions*’, hep-th/9704158
- [53] P. Horava, ‘*Matrix Theory and Heterotic Strings on Tori*’, hep-th/9705055;
D. Kabat and S.-J. Rey, ‘*Wilson Lines and T-Duality in Heterotic M(atric) Theory*’, hep-th/9707099.
- [54] S. Govindarajan, ‘*Heterotic M(atric) theory at generic points in Narain moduli space*’, hep-th/9707164.

- [55] M. Berkooz, and M. Rozali, ‘*String Dualities from Matrix Theory*’, hep-th/9705175.
- [56] D. Morrison and C. Vafa, ‘*Compactifications of F-Theory on Calabi–Yau Threefolds – I*’, Nucl.Phys. **B473** (1996) 74, hep-th/9602114;
D. Morrison and C. Vafa, ‘*Compactifications of F-Theory on Calabi–Yau Threefolds – II*’, Nucl. Phys. **B476** (1996) 437, hep-th/9603161.
- [57] E. G. Gimon and C. V. Johnson, ‘*Multiple Realisations of $\mathcal{N}=1$ Vacua in Six Dimensions*’, Nucl. Phys. **B479** (1996), 285, hep-th/9606176
- [58] E. Gimon, private communication. See also ref.[59].
- [59] P. Berglund and E. Gimon, to appear.